

Revisions of Australian ground-hunting spiders: I. *Amauropelma* gen. nov. (Araneomorphae: Ctenidae)

Robert J. Raven¹, Kylie Stumkat¹ and Michael R. Gray²

¹ Queensland Museum, PO Box 3300, South Brisbane, Queensland 4101, Australia

² Australian Museum, College St, Sydney South, New South Wales 2000, Australia

Abstract – Revisions of Australian ground-hunting spiders: I. *Amauropelma* gen. nov. (Araneomorphae: Ctenidae). A new genus *Amauropelma* is described from north-eastern Australia and includes 16 new species: *Amauropelma anzses*, *Amauropelma bluewater*, *Amauropelma claudie*, *Amauropelma gayundah*, *Amauropelma gordon*, *Amauropelma hasenpuschi*, *Amauropelma leo*, *Amauropelma mcilwraith*, *Amauropelma monteithi*, *Amauropelma mossman*, *Amauropelma pineck*, *Amauropelma rifleck*, *Amauropelma torbjorni*, *Amauropelma trueloves* (type species), *Amauropelma undara*, and *Amauropelma wallaman*. The characters of the genus challenge existing family diagnoses of the Miturgidae and Ctenidae.

INTRODUCTION

Araneomorph spider (and to a lesser extent mygalomorph) systematics are currently in a severe state of flux. Platnick, Ovtsharenko and Catley (e.g. Platnick, 2000) are revising the diverse Australian Gnaphosoidea, Diana Silva is revising the Ctenidae, Griswold and others are examining relationships of the Lycosoidea, and Raven and Stumkat are revising the Australian Clubionidae, Miturgidae, Ctenidae, and Zoridae. Family boundaries are changing but stability is coming. The genus described here is one such example. The ambiguity of its placement was nominally resolved only when Dr Diana Silva, American Museum of Natural History, on examining the genus, unequivocally declared it to be a ctenid.

Ctenids are cursorial spiders with two claws, claw tufts, and two recurved eye rows which when viewed from the front it can be seen that the ALE are set high near posterior eyes (e.g., Dippenaar-Schoeman and Jocqué, 1997). They are typically tropical spiders of the rainforest, heathland, open forest and desert. In most collections, they are still quite rare. The Australian Ctenidae at present include *Ctenus agroecoides* (Thorell, 1881) from northern Queensland, the blind *Janusia muiri* Gray, 1973 from caves in Western Australia, and *Leptoctenus agalenoides* L. Koch, 1878 from Rockhampton, Mackay, mid-eastern Queensland. No members of the family were listed by Rainbow (1911). The spiders described here are small and hunt freely in the rainforest litter of north-eastern Queensland. They have not been found associated with webs or carrying an egg sac. Reports on surveys of the fauna of north-eastern Queensland placed them either in the Miturgidae or Ctenidae

(Davies, 1976, 1977), depending on the anterior eye row curvature. When those taxa are examined together it is clear that the eye differences in this group are interspecific. Hence, the diagnosis of the Ctenidae is challenged and the boundary with the Miturgidae is diffused.

The concept of the Miturgidae has been ambiguous, partially addressed but not resolved by Griswold (1991, 1993, 1994). The family (sensu Lehtinen, 1967 and Griswold, 1993, 1993) included claw-tufted spiders with two recurved eye rows (e.g. *Diaprogapta* Simon, 1909), although in most genera the back eye row is straight or slightly procurved. The boundary with the Ctenidae seemed clear: ctenids have the smaller ALE set high above AME, near the PLE, forming an eye formula of 2.4.2. (The zorid genus *Elassoctenus* has ALE set higher but they are closer to the PME than the PLE and do not form the 2.4.2 formula.) The inclusion then of some undescribed species of *Amauropelma* gen. nov. in the Miturgidae and others in the Ctenidae (e.g. Davies, 1976, 1977) indicates that the eyes of different species fit different families.

The species described here are unequivocally one genus. Males lack tibial cracks and cymbial scopula and the tapetum is grate-shaped. Hence, the genus cladistically rises above the Tengellidae and Zoropsidae (as grouped by Griswold, 1993). Females have lateral teeth on the epigyne and, along with the curvature of the eye rows, the Ctenidae are clearly their closest relatives although cladistically *Amauropelma* may be the sister group of most other ctenids.

This is the first of a series of papers revising the family in Australia.

MATERIALS AND METHODS

All material of *Amauropelma* was collected in north-eastern Queensland. The left palp is described with the ventral face dorsal. Eye group size is measured through the centre of the group and compared to the carapace width on that line. Abbreviations and methods are standard for the Araneae and explained in Raven (1994). Hair terminology (plumose, feathery) follows Lehtinen (1967). Spination on metatarsi III and IV test the repeatability of spine counts because the boundary of lateral versus ventral spines is often vague. As before (e.g. Raven, 1994), a spine is considered lateral if the base is visible when viewed from above. Measurements, if not otherwise indicated, are in millimetres. Colour is taken from alcohol-preserved specimens unless noted. The median apophysis is cup- or ladle-shaped. The prolateral portion is termed the handle (of the ladle) and is roughly taken from the edge of the lumen. Often the lumen is not visible in ventral view but only from distal axial view or lateral view. The epigyne consists of a plate with two lateral lobes, here termed 'wings', and the lateral teeth are also described as 'horns'. The dimensions of the epigyne are taken from the widest points and length is measured from the plates, not including lateral teeth, and not from the central dimension (see Figure 14g).

Authorship. Statements concerning relationships are those solely of the senior author.

Acronyms: AMNH, American Museum of Natural History, New York; AMS, Australian Museum, Sydney; ANIC, Australian National Insect Collection, Canberra; ANZSES, Australian New Zealand Scientific Exploration Society; QM, Queensland Museum, Brisbane; SAM, South Australian Museum, Adelaide; WAM, Western Australian Museum, Perth.

SYSTEMATICS

Family Ctenidae

Amauropelma gen. nov.

Type Species

Amauropelma trueloves sp. nov.

Diagnosis

Ecribellate lycosoid with two claws, well-developed claw tufts, eight eyes in two clearly recurved rows, or blind, eyes of similar size, ALE sometimes set back near PLE, tapetum (of PME) grate-shaped; lateral spinnerets all short, conical, with short apical segment; ALS clearly separated; colulus absent; PLS of females cylindrical; tarsal organ (or rod) on legs III, IV set at basal 1/8–1/6;



Figure 1 *Amauropelma trueloves* sp. nov., female, habitus.

more distal on legs I, II. Tibiae I, II with 4–5 pairs of strong spines ventrally; 3 strong pairs ventrally on metatarsi I, II; trochanters deeply notched. Retrocoxal hymen present on leg I; pre-distal tarsal fracture absent. Tarsi with adpressed trichobothria (Figures 10b, 16a). Males without tibial crack on legs; processes retrolaterally on distal patella and tibia of palp; cymbium apically coniform, without scopula dorsally; retrobasal and probasal processes on cymbium of most species; bulb with subtegular-tegular interlocking lobes; tegulum large with cup-shaped median apophysis, hyaline (filmy) conductor; embolus a large hook-shaped plate, directed clockwise, with 2–4 deep ventral ridges narrowing distally; sperm duct opens on back of plate. Epigyne a plate with soft lateral teeth arising



Figure 2 *Amauropelma mossman* sp. nov., male, carapace and abdomen, dorsal view. Scale line, 1 mm.

from its dorsal surface; spermathecae simple. Tracheal spiracle near spinnerets, indistinct.

Description

Cephalothorax pear-shaped; caput broad, low; uniform cover of fine hairs, none feathery or plumose; short fovea. Blind or eight small, similarly-sized eyes, sessile or normal on raised black tubercle in two very recurved rows; rows are 2.2.2.2, 2.4.2 or 4.4. Median eyes set close together, ca. 1 diameter apart; MOQ longer than wide; group narrower in front than behind, occupying 0.5–0.7 of head-width. Eye region gradually sloping; from in front, AME round set on common low tubercle;

ALE oval set high but not beside PME. Chelicerae partially porrect, parallel-sided, not apically coniform, with small low boss. Fangs without ridges or grooves, not elongate; tooth margins short, diagonal with 4 large and 1–2 smaller teeth on retromargin, 2–4 on promargin. Maxillae long, ca. twice labium length; anterior ectal face truncate, outer edge broadly rounded; lunate depression on midectal face, basally truncate; serrula in long curving line. Labium clearly longer than wide, no strong grooves/ depressions, apically truncate, not convergent. Sternum shield-shaped, margins with deep scallops for anterior inner corner of coxa II, III; posteriorly blunt.




Figure 3 *Amauropelma*, females, scanning electron micrograph: a, *A. anzses* sp. nov., tarsal organ (t.o.) leg I; b, *A. trueloves* sp. nov., tarsi I showing position of tarsal organ (t.o) and enlarged dorsal setae; c, d, *A. rifleck* sp. nov., epigyne, dorsal face; e, *A. mcilwraith*, lateral palp and bulb showing interlocking lobes (i.l); f, g, *A. hasenpuschi*, female, leg I, showing tarsal organ and prostrate trichobothria (f), and claws and tufts (g).

Legs. Formula 4123 (in all measured females save *A. hasenpuschi* = 4132, most males) or 4132 (males of *A. trueloves*, *A. bluewater*, *A. gayundah*, *A. rifleck*, *A. wallaman*). Coxae all similar in shape, basally with small but distinct precoxal sclerites (inner triangular extensions), posterior more distinct than anterior; large distinct retrocoxal hymen basally on I, no opposed setae evident on prolateral II. All trochanters deeply notched. Hairs all simple. Scopula entirely absent except in ♀ *A. undara* and ♂ *A. anzsies*. Pilosity generally light. Spines: all strong; strong paired spines ventrally on tibia (4–5 pairs) and metatarsi (3 pairs) I, II, not or barely overlapping on tibiae; long, strong and overlapping on metatarsi; weakest on femora, absent on patellae I, II; proventral spine on femora absent or weak on femora I, III, and sometimes IV. Basal pair of spines on ventral tibiae I, II well-separated from base. Common spine pattern: I, II: pa, 0; ti v2.2.2.2.2; me v2.2.2. III, IV: ti v2.2.2. Preening combs absent. Claw tufts dense, narrow but set well below long claws. Two elongate claws with few small teeth basally; claws relatively longer on III, IV. Cuticle surface more or less smooth. Trichobothria in 2 rows on tarsi, irregular row on metatarsi, diffused band on tibiae. Tarsal trichobothria of two types: adpressed and erect (Figure 10b). Adpressed with more widely spaced microfilaments, base and aperture smaller than for erect trichobothria; base roughly triangular and aperture not on raised dome. Erect trichobothria larger; bases semicircular and aperture on raised dome often with distal transverse ridge; both types have 2–4 transverse ridges which may or may not be continuous across base. Trichobothria not in lengthening row on tarsi. Tarsal organ low, set in basal 1/6th of tarsi, ovate on III, IV, mid-distal on I, II, delimited by shallow encircling groove and distal acentric ovoid aperture; aperture very large in *A. undarain*; in *A. anzsies* (Figure 10c) and *A. mossman*, tarsal organ is distinct raised cup with large aperture dominating distal vertical face; in *A. monteithi*, tarsal organ an elongate inclined rod with lobular apex with large aperture (Figure 23b). Tarsi dorsally with two irregular lines of very large, thick setae with enlarged bases (Figure 3b).

Female palp: claw dentate, long, without tuft; pair of spines ventrally on tarsi predistally, also on cymbium (some species) and also probasally on tibiae.

Spinnerets very small, short; colulus absent; ALS about twice size of PLS; PMS (female) small, cylindrical, apical segments domed.

Male palp: patella with coniform process off each corner of patellar fold; tibia with retrodorsal conical mound bearing bifurcate apophysis. Cymbium boat or leaf-shaped with strong probasal lobe and retrobasal lobe or ridge, bent strongly at base of apical cone; weak scopula sometimes dorsally on

cymbium; cymbium not deep, without retrobasal groove; apically conical, two spines ventro-apically. Subtegular-tegular locking lobes present; tegulum C-shaped; embolus a large curving plate, directed clockwise, with 2–4 deep ventral ridges narrowing distally; sperm duct opens on back of plate (Figures 6c, 33d,e); median apophysis arising from large, central, pallid, unsclerotised region, ladle-like; conductor large, distal, membranous.

Epigyne a large trilobed plate with 2 horn-like processes (=teeth) below and behind lateral lobes; lateral and anterior plate margins rolled under (Figure 3c, d); anterior septum present only in *A. claudie*; simple spermathecae with anterior copulatory fossae behind plate; insemination duct follows curve of lateral lobe and enlarges distally to spermathecae; fertilisation duct posterior horn-like. In *A. undara*, an epigynal plug was present on one side of the epigyne under the anterior outer quadrant of the plate.

Tracheae (in *A. rifleck*) are two very short tubes limited to posterior 1/6 of abdomen and arising from a common lumen. Tracheal spiracle short, barely evident.

Remarks

Amauropelma shares with *Janusia* the elongate form of the tarsal claws, the shape of the carapace, leg spination, spinneret configuration, and leg pilosity, and absence of leg scopula. *Amauropelma* differs from *Janusia* in that the tarsal organ or rod is basal rather than subdistal and the opening of the organ is circular not triangular. Also, in *Amauropelma*, the epigynal plate is wide with two 'wings' rather than long; in *Amauropelma*, the copulatory fossae are in front of the leading edge of the plate, whereas in *Janusia* they are lateral. In *Janusia*, the lateral teeth are processes extending from the lateral epigynal sides whereas in *Amauropelma* they begin internally just behind the copulatory fossae and extend back and out. Also, in *Janusia*, the maxillae are apically divergent (according to Gray, 1973, figure 2) whereas in *Amauropelma* they are parallel-sided.

Unnamed ctenids with eyes from caves in Western Australia strongly resemble *Janusia*. In these species, the lateral epigynal teeth are produced from the distal edge of the epigyne, the relative eye sizes are different (ALE = AME < PME = PLE) and the ALE are closer to the PLE than PME but not on a common mound.

Amauropelma also shares similarities with the South African *Phanotea* Simon in the form of the male palp which also has a scooped median apophysis and the embolus apically divided into two divisions (see Griswold, 1994). Unlike *Phanotea*, *Amauropelma* has claw tufts and lacks both a third claw and the tibial crack in males; most species also lack leg scopula. Moreover, unlike *Phanotea*,




Figure 4 *Amauropelma* records, north-eastern Queensland: inset, Cape York.

Amauropelma has a patellar spur on the male palp, a pro- and retrobasal cymbial process, and a conically tapered cymbium.

Distribution and Habitat

Most species are found in lowland rainforests from Iron Range (200 km south of the tip of Cape York) to just north of Townsville, north-eastern Queensland; one species lacking eyes has also been taken from caves west of Townsville.

Phenology

Males of *Amauropelma* have been taken throughout the year in *A. trueloves*, in which sampling has been year-long, five males were taken in winter (June). Hence, the generally accepted notion that araneomorphs are summer maturing must be tested with each genus.

Relationships

Relationships with taxa outside of Australia are complex principally because hypotheses about relationships among lycosoid genera are in a state of flux. On an ABRIS funded visit to European and American Museums, RJR examined diverse genera while surveying the retrocoxal hymen. Some data presented here derive from that study.

Amauropelma was added to the data set of Griswold (1993) along with the retrocoxal hymen, the predistal tarsal fracture (see Raven, 1998) and

the (dorsoventral) depth of the cymbium. Using the same character coding and Hennig86 commands as Griswold (1993), one tree was found. The tree placed *Amauropelma* in the ctenoid clade as the sister group of *Ctenus* and *Phoneutria*. This coincides with conclusions formed following discussions with Diana Silva: that *Amauropelmalain* is a ctenid. However, the plesiomorphic condition of the eyes in *Amauropelma* is a mildly recurved front row (with ALE beside AME, and not beside PME as in ctenids); hence, its inclusion in the Ctenidae completely compromises the traditional diagnostic character of the group.

Amauropelma seems most closely related to the African-Asian genus *Thoriosia* Simon but differs in the presence of a patellar spur on the male palp, the basal tarsal organ and in the similar sizes of the eyes; in *Thoriosia*, the AME are much smaller than the PME (Benoit, 1976). This relationship was suggested by Silva who will explore that question more deeply.

Other similar taxa

Three other genera show some indication of a relationship which *Amauropelma*. The New Zealand amaurobiid *Maloides cavernicola* (Forster and Wilton, 1973, holotype in New Zealand Arthropod Collection, examined) is known only from a single, three-clawed, female. However, in the eye group size (reduced), small eyes, two recurved (as figured, Figure 947, not procurved as described) eye rows, epigyne with lateral teeth and median plate, and simple spermathecae, the genus is similar to *Amauropelma*.

The Chilean 'amaurobiid' *Emmenomma* has 3 claws (no tufts), a scooped median apophysis and lateral teeth on the epigyne. It may also belong in this group. Third, Lehtinen (1967: 317) commented on presumed adaptations to cave living of some species of *Phanotea* – length of legs and trichobothria, reduced colour pattern, size of eyes and lengthening of the straight distal part of the paired tarsal claws – all found in *Amauropelma*.

Cladistics of Species

Within the genus, two characters show phylogenetically interpretable differences: the curvature of the front eye row and the development of the male patellar apophyses on the palp.

The front row eye curvature varies from almost straight from above to clearly curved to the extent that the ALE are decidedly higher than the AME and approach the PME (see *A. claudie*, Figure 14f). The outgroup condition is a more or less straight line and the ALE set beside the AME. Hence, clearly, the plesiomorphic condition of the front row curvature is more or less straight and the genus should not qualify on that character to be included in the Ctenidae.

The palpal patella has a retrolateral invagination in many araneomorph genera. In *Amauropolma* males, the distal corners of the invagination form processes varying from small cones (*A. bluewater*) to at least one large tube (*A. gordon*). The outgroup condition (absence) suggests that the character is a synapomorphy of the genus.

Two species-groups are recognisable by a combination of characters: in the *leo* group, the eyes and overall eye group are small, the carapace lacks pattern and the spiders are larger; in the *trueloves* group, the eyes are larger and the group is wider relative to the head width, the spiders are smaller and the carapace has distinct lateral dark bands.

Morphology

Tarsal organ and rod. Typically, the tarsal organ is distal on the tarsi in spiders. When in the form of a rod in mygalomorphs (*Ixamatus*, *Xamiatus*; Raven, 1981) it is also distal. Araneomorph genera with a tarsal rod include the gradungulids *Tarlina* and the amaurobiids *Otira* and *Strenosoma* (e.g. Davies, 1986). Davies (1986: 240) noted that in *Otira* the rod is distal in New Zealand species and basal (ca. 0.25 length) in Australian species. The homology of the diverse rods is not presumed but simply the elevation of the receptive component off the tarsal surface. In *A. monteithi*, the rod is in the same location as the tarsal organ in other species and also there is no other tarsal organ; hence, in this case the rod and organ are presumed homologues. On all legs, the rod is at the basal 1/5.

The retrocoxal hymen (first reported by Raven, 1998) is readily observed by light microscopy on a medium-sized spider of the genus *Cheiracanthium* in which it is evident as a pallid unsclerotised circle on the ental half of the retrolateral face of coxa I. The predistal tarsal fracture (first reported by Raven, 1998) when present, occurs on all leg tarsi and is sometimes more easily seen in lateral view by light microscopy as a pallid diagonal zone predistally on the tarsus.

The embolus is unlike any thus far seen in Australian miturgids, zorids, clubionids, cycloctenids, pisaurids, or other ctenids. It consists of a broad flattened plate with several high curving ridges that distally converge and presumably 'lock' into the epigynal teeth or horns. The physical sizes are similar, at least for the distal portion of the embolus. The base of an epigynal horn is about 90µm wide and 200µm long; the embolus tip is about 50µm long and 20 µm wide. The opening of the sperm duct is discernible only under an SEM (Figures 6c, 33d, e) with the embolic plate so inclined as to allow examination of the dorsal apex of the plate. Prolaterally, the embolus is bounded by a curved ring, deeply concave externally, which probasally has an interlocking lobe with the tegulum (Figure 3e).

Species Included

Amauropolma anzses sp. nov., *A. bluewater* sp. nov., *A. claudie* sp. nov., *A. gayundah* sp. nov., *A. gordon* sp. nov., *A. hasenpuschi* sp. nov., *A. leo* sp. nov., *A. mcilwraith* sp. nov., *A. monteithi* sp. nov., *A. mossman* sp. nov., *A. pineck* sp. nov., *A. rifleck* sp. nov., *A. torbjorni* sp. nov., *A. trueloves* sp. nov., *A. undara* sp. nov. and *A. wallaman* sp. nov.

Etymology

From the leg pattern which is reminiscent of the amaurobiid basic pattern; the gender is neuter.

Key to species of *Amauropolma*

Males (males of *A. claudie* unknown)

1. Eyes absent *A. undara*
Eyes present 2
2. Long tarsal rod present basally (Figure 23b)
..... *A. monteithi*
Tarsal rod absent or very low 3
3. Patellar apophysis long, vermiform (Figure 14c,
15b) *A. gordon*
Patellar apophysis short, coniform if present .. 4
4. Tarsal organ posteriorly raised with vertical
distal face on mid-tarsi on I, basal 1.4 on II-IV
(Figure 3a) *A. anzses*
Tarsal organ low, indistinct 5
5. Median apophysis subcircular with dominant
cavity (Figure 12a) *A. bluewater*
Median apophysis clearly longer than wide 6
6. Patellar apophysis a prominent cone (Figure
31c) *A. torbjorni*
Patellar apophysis a small sometimes bifid
cone, if present 7
7. Retrolateral paracymbial lobe distinctly
produced (Figure 6a); cavity of median
apophysis confined to apical one-third; large
serrate keel on base of embolic plate (Figure
6d) *A. trueloves*
Retrolateral paracymbial lobe absent or thick
but not well produced (Figures 18a, 24a) 8
8. Tibial apophysis unequally bifid or single with
small subdistal tooth (Figure 27b) *A. pineck*
Tibial apophysis equally bifid (Figures 29b, d) .
..... 9
9. Patellar apophysis is one fused bifid process
(Figure 29c) *A. rifleck*
Patellar apophysis is clearly two separate
cones, if present (Figure 24c) 10
10. Median apophysis without lumen (Figures 13a,
27f) *A. gayundah*

- Medan apophysis clearly with lumen (e.g. Figure 24a) 11
11. Embolic plate with thumb-like lobe below tip (Figure 20b) *A. mcilwraith*
 Embolic plate without thumb-like lobe below tip 12
12. Tibial apophysis deeply divided for apical 2/3 (Figure 34d) *A. wallaman*
 Tibial apophysis shallowly divided in apical 1/3 (Figure 18b) 3
13. Median apophysis with dominant (prolateral) 'handle' (Figure 24a, b); embolic plate with low basal extra keel *A. mossman*
 Median apophysis with dominant (retrolateral) lumen; basal embolic plate without extra keel (Figure 18c) *A. hasenpuschi*
- Females (females of *A. gayundah* unknown)**
1. Eyes absent *A. undara*
 Eyes present 2
2. Long tarsal rod present (Figure 23b)
 *A. monteithi*
 Tarsal rod absent or very low 3
3. Tarsal organ posteriorly raised with vertical distal face on mid-tarsi on I, basal 1.4 on II-IV (Figure 10c); scopula present on tarsi I, II
 *A. anzsies*
 Tarsal organ low, indistinct; scopula absent on tarsi 4
4. Epigynal plate subquadrate (Figures 11c, 17f) ..
 5
 Epigynal plate clearly wider than long (Figure 5f) 6
5. From above, ALE about midway between PLE and AME (Figure 17d) *A. hasenpuschi*
 From above, ALE clearly closer to PLE than ALE (Figure 11b) *A. bluewater*
6. Distal epigynal plate with concave cavity to lateral lobes (Figure 5f) 7
 Distal epigynal plate with clearly convex lateral curve from distal edge to lateral wings (Figure 14d) *A. gordon*
7. Lateral horns set wide of and orthogonal to sagittal plane and extend well beyond lateral wings (Figure 14g) *A. claudie*
 Lateral horns set close to lateral wings or hardly wider (Figure 5f) 8
8. Eye group small (Figure 19b) and AME clearly larger than PME *A. leo*
 Eye group large, normal (Figure 5a); AME and PME similar sizes (Figure 5d) 9
9. From above, front edge of ALE in same line as those of PME or behind them; PME clearly larger than AME (Figure 25a) 10
 From above, front edge of ALE clearly in front of those of PME; PME equal or subequal to AME (Figure 30a) 13
10. Tibiae I, II with 5 pairs of spines ventrally 11
 Tibiae I, II with only 4 pairs of spines ventrally *A. pineck*
11. Overall epigyne only ca. 1.5–1.6 times wider than long (Figure 25c) 12
 Overall epigyne wider, ca. 1.9 times wider than long (Figure 5f); north of Daintree River and south of Bloomfield River *A. trueloves*
12. PME clearly bigger than ALE (Figure 25a); just south of Daintree River *A. mossman*
 All eyes of similar size; just north-east of Townsville *A. wallaman*
13. Overall epigyne wide ca. 2.1 times wider than long with slender lateral horns (Figure 28b); Ellis Beach region *A. rifleck*
 Overall epigyne longer ca. 1.7 times wider than long with fleshy lateral horns (Figure 30f); Mt Cook region *A. torbjorni*

Amauropelma trueloves

Raven and Stumkat, sp. nov.

Figures 1, 3b, 4–7, Table 1

Material Examined

Holotype

♂, Pilgrim Sands, Cape Tribulation, Queensland, Australia, 16°00'S, 145°25'E, 24–29 August 1988, R.J. Raven (QM S24194).

Allotype

Australia: Queensland: 1 ♀, Cape Tribulation, 2 km WNW (Site 2), 16°04'S, 145°27'E, pitfall, 23 September–7 October 1982, G. Monteith, D. Yeates, G. Thompson (QM S31355).

Paratypes

Australia: Queensland: 1 ♀, 24–29 August 1988, R.J. Raven (QM S32917); 1 ♀, 27 March–1 May 1996, P. Zborowski (QM S39107); 1 ♂, sieved litter, 13 October 1980, G. Monteith (QM S32888); 1 ♂, sieved litter, 14 October 1980, G. Monteith (QM S32885); 1 ♀, sieved litter, 23 September 1982, G. Monteith, D. Yeates, G. Thompson (QM S32886); 1 ♀, sieved litter, 2 October 1982, G. Monteith, D. Yeates, G. Thompson (QM S32887); 3 ♀, pitfall, 31 May–28 June 1996, P. Zborowski (QM S41784); 1 ♀, 5 December 1995–4 January 1996, L. Umbach (QM S39113); 1 ♂, pitfall, 31 May–28 June 1996, P. Zborowski (QM S39112); 2 ♂, pitfall, 31 May–28




Figure 5 *Amauropelma trueloves* sp. nov.: holotype male, a–d: a, carapace and abdomen, dorsal view; b, leg I, prolateral view; c, sternum, maxillae, labium and coxae; d, eyes, dorsal view. e–i, female: e–h, allotype; e, abdomen, dorsal view; f, epigyne, external, ventral view; g, carapace and chelicerae; h, eyes, frontal view; i, epigyne (QMS41785, dorsal view. Scale lines = 1mm, except f, i, 0.5mm.

June 1996, P. Zborowski (QM S39109); 1 ♀, pitfall, 1–28 February 1996, L. Umback (QM S39116); 1 ♀, sieved litter, January 1983, G. Monteith (QM S26902); 1 ♂, flight intercept trap, 1–28 February 1996, L. Umback (QM S39114); 1 ♂, flight intercept trap, 31 May–28 June 1996, P. Zborowski (QM S41782); 1 ♀, pitfall, 31 May–28 July 1996, P. Zborowski (QM S39105); 1 ♀, pitfall, 31 May–28 July 1996, P. Zborowski (QM S39106); 1 ♀, pitfall, 31 May–28 June 1996, P. Zborowski (QM S39110); 1 ♀, pitfall, 5 December 1995–4 January 1996, L. Umback (QM S39118); 3 ♀, pitfall, 27 March 1996–1 May 1996, P. Zborowski (QM S41785); 1 ♂, pitfall, 31 May–28 June 1996, P. Zborowski (QM S39111)); 1 ♀, pitfall, 1–28 February 1996, L. Umback (QM S39117); 1 ♂, 1 ♀, pitfall, 1–28 February 1996, L.

Umback (QM S41783); 2 ♂, sieved litter, 23 September–7 October 1982, G. Monteith, D. Yeates, G. Thompson (QM S32884); 1 ♀, Noah Head, S. Cape Tribulation, 16°07'S, 145°27'E, sieved litter, 16 October 1980, G. Monteith (QM S26904); 2 ♂, pitfall, 29 November 1992–17 April 1993, R., J. and S. Raven, P. and E. Lawless (QM S19769); 2 ♀, pitfall, 21 July–29 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QM S24038); 13 ♂, 3 ♀, pitfall, 5 November 1991–20 July 1992, R. Raven, P. Lawless, M. Shaw (QM S24863, WAM T42600, AMNH, ANIC). All from rainforest at Cape Tribulation.

Diagnosis

Differs from *A. monteithi* in lacking a tarsal rod, having larger raised eyes and more pattern and




Figure 6 *Amauropelma trueloves* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia; b, tibial apophysis, ventral view; c, embolus tip showing spermatic outlet, axial view; d, embolic plate and conductor, ventral view; e, median apophysis, ventral view; f, patellar apophyses, retrolateral view.

from *A. pineck* in have 5 pairs of spines ventrally on tibiae I, II.

Description

Holotype male (QM S24194)

Carapace 2.46 long, 2.00 wide. Abdomen 2.07, 1.38 wide. *Colour*. Carapace and chelicerae orange with shadows along edge of carapace, beside fovea, and as slightly dark rings on legs as in ♀. Abdomen darker with pallid medial region anteriorly, broken

open brown areas posteriorly; ventrally almost entirely pallid with darker spots laterally. Eyes all surrounded by black pigment. *Carapace*. Fovea begins at widest point. *Eyes*. AME:ALE:PME:PLE, 7:7:9:8. *Spines*. As for genus except as noted. I: fe pv1p1d3r1; ti p1. II: fe pv1p1d3r1; pa 0; ti p1; me p1. III: fe pv1p2d3r3; pa r1; ti p2d3r2; me p4r4v2.2.4 short pair medially. IV: fe pv1p2d3r2; pa r1; ti p2d3r2; me p5r5v2.2.2 short pair medially. *Palp*: fe p1d3, pa p1w, ti p2w, cymbium 2 distoventral. *Legs*. Scopula absent. *Palp*. Palpal patella with two clearly

Table 1 Leg measurements of *Amauropelma trueloves*.

Holotype male (QMS24194).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.00	1.69	1.69	2.23	0.92
Patella	0.92	0.77	0.85	1.00	0.31
Tibia	1.92	1.46	1.23	2.00	0.38
Metatarsus	1.46	1.31	1.38	1.92	
Tarsus	0.92	0.85	1.00	1.61	1.00
Total	7.22	6.08	6.15	8.76	2.61
Allotype female (QMS31355).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.92	1.73	1.61	2.23	0.92
Patella	0.85	1.08	0.85	0.85	0.38
Tibia	1.69	1.38	1.15	1.85	0.46
Metatarsus	1.23	1.23	1.31	2.15	
Tarsus	0.65	0.73	0.92	1.15	0.69
Total	6.34	6.15	5.84	8.23	2.45





Figure 7 *Amauropelma trueloves* sp. nov. female, scanning electron micrographs: a, distal tarsus, paired claws and claw tufts; b, tarsal organ and trichobothium.

equally sized divided cones on retrodistal corner; retrobasal cymbial lobe rounded, blunt bur distinctly produced; embolic plate short, beak-like. Median apophysis long, ladle-shaped with small lumen and long broad handle. *Spinnerets*. All very short or small, atrophied; PMS=PLS <0.5x ALS; all with domed tips. Slight but evident space between ALS.

Allotype female (QM S31355)

Carapace 2.54 long, 2.08 wide. Abdomen 3.30, 1.92 wide.

Colour. More boldly marked than *A. monteithi*. Carapace yellow orange with narrow marginal bands on each side stopping before anterior constriction, wider dark band begins at AME across to anterior carapace edge and back beside fovea. A wide dark band medially down chelicerae; abdomen dark grey with large pallid medial area anterior and 3 paired patches behind and large patches laterally. Legs strongly banded; 3 bands on femora only distal unbroken dorsally by spines. One band on patella and two on each of tibia and metatarsus. Ventral abdomen pallid with roughly large paired areas of dark. Sternum, maxillae and labium yellow orange, chelicerae darker; irregular shadows medially on sternum. Eyes with dark bases. *Carapace*. Almost glabrous save for fine black hairs (and sheen). Paired setae along clypeal margin, single bristle between AME. *Eyes*. AME:ALE:PME:PLE, 7:8:9:9. Eyes seem relatively larger cf male. *Chelicerae*, maxillae and labium similar to male. *Spines*. As for genus except as noted. I: fe pv1p1d3. II: fe p1d3. III: fe pv1p2d3r2; pa r1; ti p2d3r2; me p4r2v2.2.2. IV: fe pv1p2d3r1; pa r1; ti p2d2r2; me p4r4v2.2.2. *Palp*: fe p1d3; pa p1; ti p2; ta p3. *Legs*. Scopula absent. Trochanters deeply notched (notch as long as wide on IV less so on I but still deep). Sheen also present. *Claws*. 2, noticeably long and thin, 3–4 short teeth medially. Tufts small, below claw distinct. *Epigyne*. Externally a broad plate with two deeply encased broad lateral lobes and long curved horns. Copulatory ducts in horns.

Variation

Intensity of pattern on carapace, legs and abdomen of males varies from pallid to dark.

Distribution and Habitat

Known only from lowland rainforest near Cape Tribulation, north of Cairns, north-eastern Queensland.

Etymology

A noun in apposition from a local name for the area.

Amauropelma anzses Raven and Stumkat, sp. nov.

Figures 3a, 4, 8–10, Table 2

Material Examined

Holotype

♂, Mt Finnigan, site 1, Queensland, Australia, 15°47'S, 145°16'E, pitfall, 4 December 1990–17 January 1991, QLD Museum and ANZSES (QM S31360).

Paratypes

Australia: Queensland: 2 ♂, same data as holotype (QM S31359); 2 ♂, same data as holotype except site 3 (QM S32918); 1 ♂, same data as holotype except site 4 (QM S32919); 1 ♂, Big Tableland, 15°43'S, 145°16'E, flight intercept trap, 20 December 1990–8 January 1991, ANZSES expedition (QM S32914).

Diagnosis

Differs from all other species in having leg scopula in males and in having a distinctly raised (but not a rod) tarsal organ.

Description

Holotype male (QM S31360)

Carapace 4.54 long, 3.77 wide. Abdomen 4.07, 2.69 wide.

Colour. Entirely yellowish, dark fine lines along strial margins on carapace. **Eyes.** AME:ALE: PME:PLE, 10:8:6:6. AME clearly the largest. **Chelicerae.** Promargin with 2 teeth; retromargin with




Figure 8 *Amauropelma anzses* sp. nov.: holotype male: a, carapace and abdomen, dorsal view; b, eyes, dorsal view; c, eyes, front view; d, spinnerets, ventral view. Scale line = 2mm for a, 1mm for b–d.

Table 2 Leg measurements of *Amauropelma anzses*.

Holotype male (QMS31360).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	4.08	3.69	3.31	4.31	1.77
Patella	1.92	1.92	1.46	1.69	0.61
Tibia	4.08	3.31	2.92	4.08	0.77
Metatarsus	3.69	2.92	3.15	4.61	—
Tarsus	2.07	1.46	1.38	2.38	1.46
Total	15.84	13.30	12.22	17.07	4.61

4. **Sternum.** Posterior corners almost flat, others distinct but broad and mid-distal. **Spines.** As for genus except as noted. I: fe pv1 strong p2d3r2; ti p1r1; me p2r1. II: fe pv1p2d3r2; ti p1r1; me p3r2. III: fe pv1p2d3r4; pa r1; ti p2d3r2; me p2+2l r3+2l v2.2.2. IV: fe p4d3 r2; pa r1; ti p2d2r2; me p2+2l r3+2p v5.2. **Palp:** fe p1d1.2; pa p1w; ti p2r1. **Cymbium,** 2 small subapical. **Legs.** Distinct raised lobe on tarsus almost medial on I, at basal 1/4 on II–IV. Trochanteral notches deep. **Scopula:** tarsi I–II, entire; IV with line of emergent setae; metatarsi I–III entire; IV tripartite. **Claws.** 3–4 short teeth on I, 1 small on IV, tufts narrow distinct. **Trichobothria.** 2 rows on tarsi, irregular row on metatarsi, diffused band on tibiae. **Palp.** Patella with two separate retrodistal cones of dissimilar height, upper cone larger. Tibia incrassate, distodorsally with dissimilarly bifurcate apex (ventrally retro-laterally), two distal extensions on tibia prolaterally interact with broadly rounded probasal cymbial lobe; retrobasal cymbial lobe a distinct process from basal cymbial flange. **Spinnerets.** Wide apart; ALS about 1.2 diameter apart, PMS long.

Female

Unknown.

Distribution and Habitat

Known only from rainforest at Mt Finnigan and Big Tableland, north-eastern Queensland.

Etymology

An acronym of Australian New Zealand Scientific Exploration Society (ANZSES) for a group that included some of the collectors.

Amauropelma bluewater

Raven and Stumkat, sp. nov.

Figures 4, 11, 12, Table 3

Material Examined

Holotype

♂, Bluewater Range, Queensland, Australia,




Figure 9 *Amauropolma anzses* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, tibial apophysis, ventral view; c, embolic plate and conductor, ventral view; d, median apophysis, ventral view; e, patellar apophyses, disto-retrolateral view.




Figure 10 *Amauropelma anzses* sp. nov., male, scanning electron micrographs: a, distal tarsus, paired claws and claw tufts; b, addressed and erect trichobothria; c, raised tarsal organ with distal aperture.

19°10'S, 146°22'E, rainforest, sieved litter, 7 December 1986, G. Monteith, G. Thompson (QM S31366).

Allotype

Australia: Queensland: ♀, same data as holotype (QM S31367).

Paratype

Australia: Queensland: 1 ♂, Mt Halifax, SE ridge,

19°17'S, 146°23'E, 21 March–10 May 1991, D. Cook (QM).

Diagnosis

Males differ from those of all other species in the small size of the patellar process on the palp and the rounded median apophysis and females from *A. hasenpuschi* in having only 4 pairs of spines ventrally on tibiae I, II.

Description

Holotype male (QM S31366)

Carapace 2.76 long, 2.07 wide. Abdomen 4.15, 2.77 wide.

Colour. Orange brown with fine dark lines along striae, legs without bands. Abdomen dorsally dark, laterally with 5 irregular ovoid areas, centrally with dark intervening triangles, laterally pallid with dark mottling; ventrally entirely unmarked. Dark shadows around eyes. *Eyes.* AME:ALE:PME:PLE, 7:6:6:6. ALE lighter colour. All of similar size. Anterior edge of ALE almost in line with front edge of PLE. *Spines.* As for genus except as noted. I: fe pv1p1d3r1; ti r1. II: fe pv1p1d3r2; ti p1r1; me p1r1. III: fe pv1p2d3r3; pa r1; ti p2d3r2; me p2+2l v2.2.2.(2.1) close. IV: fe pv1p2d3r1rv1; pa r1; ti p2d3r2; me p2+2l d1r2+2l v1.2.2.2.2. close. *Palp:* fe p1d1.2; ti p1.1; cymbium subapical 2. *Legs.* Tarsal organ at basal 1/6; tarsal trichobothria c.4–6 dorsally in basal 1/2. *Spinnerets.* Set close together. *Claws.* Tufts distinct small. STC IV and teeth medial, ca. 6 on I. *Palp* (Figure 12). Patella with two low distal processes retrolaterally; tibia with single retrodorsal apophysis deeply divided into two

Table 3 Leg measurements of *Amauropelma bluewater*.

Holotype male (QM S31366).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.08	1.92	1.76	2.24	0.80
Patella	1.04	0.92	0.80	1.00	0.48
Tibia	1.88	1.48	1.28	2.00	0.40
Metatarsus	0.92	0.92	1.52	2.24	—
Tarsus	0.84	0.80	1.32	1.32	0.88
Total	6.76	6.04	6.24	8.80	2.56

Allotype female (QMS31367).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.76	1.76	1.60	2.20	1.12
Patella	0.96	0.80	0.80	0.96	0.48
Tibia	1.60	1.40	1.20	1.92	0.40
Metatarsus	1.24	1.20	1.20	2.00	—
Tarsus	0.60	0.56	1.08	1.08	0.68
Total	6.16	5.72	5.56	8.16	2.68




Figure 11 *Amauropelma bluewater* sp. nov.: holotype male: a, carapace and abdomen, dorsal view; b, eyes, dorsal view. c, d, allotype female: c, epigyne, ventral view; d, epigyne, dorsal view. Scale line = 2mm for a, 1mm for b, 0.21mm for c, d.

strong prongs. Cymbium with large rounded retrobasal lobe. Median apophysis almost circular with large central lumen.

Allotype female (QM S31367)

Carapace 2.76 long, 2.07 wide. Abdomen 3.53, 2.39 wide.

Colour. Carapace and legs yellow brown; abdomen pattern faded. *Eyes.* AME:ALE:PME:PLE, 6:5:5:6. ALE slightly closer to PLE than AME; from above, front edge of ALE just in front of that of PME; from in front, ALE set almost above top line of AME. Back row just form 2 lines. *Chelicerae.* Promargin with 3 teeth; retromargin with 1 small basal and 4 large distal teeth. *Legs.* Tarsal organ from distal 1/3 (I) to basal 1/8–1/6(II–IV). No scopula. *Claws.* With 3 (III, IV) or 5–6 (I, II) teeth. *Spines.* I: fe pv1p2d3; ti v2.2.2.2. II: fe p1d3; pa 0; ti v2.2.2.2. III: fe pv1p2d3r2; pa r1; ti p2d3r2v2.2.2; me p1.1.2r1.1.2.v2.2.2. IV: fe p2d3r1; pa r1; ti p2d2r2v2.2.2; me p3r3v2.2.2. *Palp.* fe p1d1.2; pa p0; ti p2d1; ta p2r1. *Epigyne.* A broadly subquadrate plate with broad fleshy lateral horns closely adpressed to lateral plate.

Distribution and Habitat

Known only from rainforest at Bluewater Range,

and Mt Halifax, just north of Townsville, north-eastern Queensland.

Etymology

A noun in apposition taken from the name of the type locality.

Amauropelma claudie Raven and Stumkat, sp. nov.

Figure 4, 14g, Table 4

Material Examined

Holotype

♀, West Claudie River, Iron Range, Queensland, Australia, 12°44'S, 143°13'E, rainforest, sieved litter, 6 December 1985, G. Monteith (QM S32901).

Diagnosis

Females differ from those of all other species in the lateral horns of the epigyne extending wide of the lateral wings and being distal.

Description

Holotype female (QM S32901)

Carapace 2.28 long, 1.88 wide. Abdomen 2.40 long, 1.72 wide. Total length 5.

Colour. Carapace yellow brown with dark marginal and central broad shadow; chelicerae and legs yellow brown with bands or pattern. Abdomen with faded pattern but 3 large paired ovoid areas anteriorly and 3 transverse pallid bands posteriorly. *Eyes.* 2.4.2. PME>ALE=PLE>PME; AME set forward on common tubercle; all but ALE fully encircled by black ring, ALE with dark crest dorsally. From above, front of ALE in line with that of PME but ALE roughly equidistant from ALE and PLE; from front, front of ALE in line with top of AME. Eye group = 0.62 of headwidth; PME = 8% of headwidth. *Chelicerae.* Retromargin with 4 large and 1 small basal tooth; promargin with 3 teeth, middle largest. *Legs.* Tarsal organ at 2/5 from distal on I, II, at basal 1/6 on III, IV. Scopula absent. *Spines.* As for genus except as noted. I: fe pv1p1d2; pa v1.

Table 4 Leg measurements of *Amauropelma claudie*.

Holotype female (QM S32901).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.60	1.60	1.44	2.00	0.80
Patella	0.88	0.80	0.60	1.76	0.32
Tibia	1.40	1.20	1.00	1.60	0.48
Metatarsus	1.12	1.00	1.20	1.92	
Tarsus	0.60	0.52	1.04	1.04	0.64
Total	5.60	5.12	5.00	8.32	2.24




Figure 12 *Amauropelma bluewater* sp. nov.: male palp, ventral right, scanning electron micrographs: a, b, cymbium, bulb and distal tibia, ventral view (a), and prolateral view (b) showing extended median apophysis; c, d, patella and tibia, ventral view (c), retroventral view (d).

Table 5 Leg measurements of *Amauropelma gayundah*.

Holotype male (QMS32990).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.72	1.60	1.40	1.88	0.64
Patella	0.80	0.76	0.64	0.76	0.32
Tibia	1.44	0.84	1.80	1.60	0.36
Metatarsus	1.20	1.08	1.12	1.80	
Tarsus	0.56	0.60	0.96	0.96	0.84
Total	5.72	4.88	5.64	7.00	2.16

II: fe p1d3; me v2.2.2. III: fe p2d2r1; pa 0; ti p2d3r2; me p3r3v1.2.2. IV: fe p2d3r1; pa r1; ti p2d3r2; m3 p3r3v1.1.1.1.2. Palp: fe p1d1.2; pa p1; ti p1d1; ta p2.1 *Epigyne*. With wide plate, anteriorly indented and large lateral horns extending widely distally on plate.

Remarks

Spination of patellae I, III is unusual: unlike all other species, a spine is present ventrally on a patella (I), and only in *A. bluewater* and *A. undara* are spines absent on patella III.

Distribution and Habitat

Known only from rainforest on the West Claudie River, Cape York, Queensland.

Etymology

A noun in apposition taken from the name of the type locality.

Amauropelma gayundah Raven and Stumkat, sp. nov.

Figures 4, 13, 27f, Table 5

Material Examined

Holotype

♂, Gayundah Ck, Hinchinbrook Is, Queensland, Australia, 18°21'S, 146°13'E, rainforest, pitfalls, 8–17 November 1984, V. Davies, J. Gallon (QM S32990).

Paratype

Australia: Queensland: 1 ♂, same data as holotype but 10m altitude, 7–14 November 1984 (QM S19813).

Diagnosis

Differs from other sighted species in that the median apophysis has no lumen viewed ventrally.

Description

Holotype male (QMS 32290)

Carapace 2.12 long, 1.60 wide. Abdomen 1.72 long, 1.08 wide.

Colour. Carapace and legs light yellow. Abdomen dorsally mottled brown (paratype) with 3–4 pairs of large white spots. *Eyes*. Lateral eyes on separate tubercles; from above ALE at back edge of AME; AME:ALE:PME:PLE, 6:6:5:5. *Spines*. As for genus except as noted. I: fe pv1p1d3; ti p1r1; me p1r1. II, as for I but fe pv1p1d3r1. III: fe pv1p2d3r2; pa r1; ti p2d3r2; me p1.1.2r1.2.2v2.2.2.4. IV: fe pv1p2d4r2; pa r1; ti p2d1r2v2.1.1.1.1; me p1.1.2r1.1.2v2.2.2.4. Palp: fe p1d2; pa 0; ti p2d1; cymbium with no subapical spines. *Palp*. Patellar fold normal, broken, forms two prominent conical processes. Tibial apophysis a single deeply divided bifid retrolateral process. Cymbium with small but distinct retrobasal lobe and strongly produced probasal lobe. Median apophysis without ventrally evident lumen.

Female

Unknown.

Distribution and Habitat

Known only from rainforest at Gayundah Creek, Hinchinbrook Island, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

Amauropelma gordon Raven and Stumkat, sp. nov.

Figures 4, 14a–e, 15, 16, Table 6

Material Examined

Holotype

♂, Gordon Ck, Iron Range, Queensland, Australia, 12°43'S, 143°19'E, pitfall, 28 June 1976, V. Davies, R. Raven (QM S31356).

Table 6 Leg measurements of *Amauropelma gordon*.

Holotype male (QMS31356).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.31	1.92	1.69	2.46	1.08
Patella	0.77	1.00	0.77	0.92	0.46
Tibia	2.07	1.46	1.31	1.92	0.54
Metatarsus	2.23	1.38	1.46	2.23	
Tarsus	1.08	0.69	0.69	1.00	0.92
Total	8.46	6.45	5.92	8.53	3.00
Allotype female (QMS31357).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.00	1.92	1.85	—	0.69
Patella	1.08	0.92	0.92	—	0.38
Tibia	1.85	1.54	1.31	—	0.61
Metatarsus	1.46	1.38	1.38	—	
Tarsus	0.61	0.69	0.85	—	0.77
Total	7.00	6.45	6.31	2.45	




Figure 13 *Amauropelma gayundah* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, c, tibia and patella showing apophyses, retroventral view (b), ventral view (c).




Figure 14 *Amauropelma*. a-e, *A. gordon* sp. nov.: holotype male (a-c), allotype female (d, e): a, b, eyes, dorsal view (a), frontal view (b); palpal patella and tibia, ventral view (c); d, e, epigyne, ventral view (d), dorsal view (e); f, g, *A. claudie* sp. nov., f, eyes, dorsal view; g, epigyne, ventral view. Scale line = 2mm for a, b, 1mm for c, 0.5mm for d, e, 0.3mm for f, g.




Figure 15 *Amauropelma gordon* sp. nov.: male palp, ventral right, scanning electron micrograph, ventral view: a, cymbium, bulb and distal tibia; b, patellar apophysis; c, tibial apophysis; (d) median apophysis; all in ventral view.

Allotype

Australia: Queensland: ♀, same data as holotype (QM S31357).

Paratypes

Australia: Queensland: 2 ♂, same data as holotype (QM S31358).

Diagnosis

Males are readily distinguished from all other species by the long vermiform patellar process on the male palpal patella; females differ from those of *A. claudie* in the lateral horns being close to the plate.

Description

Holotype male (QM S31356)

Carapace 2.62 long, 2.00 wide. Abdomen 2.53, 1.69 wide.

Colour. Carapace orange with narrow shadows around eyes. Legs yellow without bands. Abdomen pallid. **Eyes.** AME:ALE:PME:PLE, 6:5:4:4, all small. **Chelicerae.** Promargin 2 teeth; retromargin 5 teeth. **Spines.** As for genus except as noted. I: fe pv1 p1d3r2; ti p1r1; me p1r1v2.2.2. II: fe pv1 p1d3r2; ti p1r1; me p1r1. III: fe pv1p2d3r3; pa p1r1; ti p2d3r2; me p5r2+2 (lower 2 strong). IV: fe p2d3r2; pa 0; ti p2d2r2; me p2+3l rp2+2l v2.2.1.2. **Palp:** fe p1d1.2; pa p1; ti p2d1. Spines on ventral and lower lateral




Figure 16 *Amauropelma gordon* sp. nov. Female, scanning electron micrographs: a, addressed and erect trichobothria on tarsi; b, distal tarsus, paired claws and claw tufts, lateral view.

tibia and metatarsi, strong and very long on metatarsi, at least half length of segment, shorter on tibiae I, II. *Spinnerets*. ALS ca. 1 diameter apart, PMS as long but thinner than PLS. *Palp*. Patella ventral joint extended into long blunt curved process. Tibia with proventral thickening and small broad process widening and then tapering near tip to blunt point, two blunt distal collars on proventral face of tibia flank broad cymbial lobe. Tegulum forming a reflected G-shape; dark groove and curved process arising off prolateral tegulum and in front of that an elaborate short hooked grooved process with translucent conductor flared dorsally. Median apophysis narrow, gourd-shaped with narrow ridge and hangs basally back over back edge of tegulum; lateral bulb very elevated. Cymbium retrobasally with slight flange forming basal ridge, no retrobasal lobe. Annuli on subtegulum; interlocking lobes evident.

Allotype female (QM S31357)

Carapace 2.62 long, 2.00 wide. Abdomen 2.84, 1.85 wide.

Colour. As for male. *Eyes*. AME:ALE:PME:PLE, 5:5:5:5. *Spines*. As for genus except as noted. I: fe pv1w p1d3r1. II: fe p1d3r1. III: fe p2d3r1; pa r1; ti p2d3r2; me p2+2l r2+2l v2.2.2. IV: fe p2d3r1; pa 0; ti p2d3r2v1.2.2; me p2+2l r2+3l v2.2.2. *Palp*: fe p1d1.2; pa p1; ti p2r2; ta p3r3. *Epigyne*. Externally broad, rounded plate without posterior projection; horns long, sinuous without obvious duct; anteriorly on (ventral) left a sclerotised horn (presumably matched by another still on female) which appears to be copulatory duct leading to small head of spermathecae and proceeding back under fold through thick walled duct to basal lobe.

Distribution and Habitat

Known only from lowland rainforest at Gordon Creek, Iron Range, Cape York Peninsula, Queensland.

Etymology

A noun in apposition taken from the name of the type locality.

Amauropelma hasenpuschi

Raven and Stumkat, sp. nov.

Figures 3f–g, 4, 17, 18, 33d, Table 7

Material Examined

Holotype

♂, South Johnstone Forestry Camp, 2 km E., Queensland, Australia, 17°36'S, 146°00'E, rainforest, pitfall, 1 December 1993–25 February 1994, J. Hasenpusch (QM S31362).

Allotype

Australia: Queensland: ♀, South Johnstone Forestry Camp, 2 km E., 17°36'S, 146°00'E, rainforest, pitfall, 1 December 1993–25 February 1994, J. Hasenpusch (QM S31363).

Paratypes

Australia: Queensland: 2 ♂, Copperlode Dam Rd (NQ 42), 16°58'S, 145°42'E, rainforest, pitfall, 27 October 1991–23 July 1992, P. Lawless, R. Raven, M. Shaw (QM S32910); 1 ♂, Graham Ra, 17°17'S, 145°57'E, rainforest, pitfall, 1 November–8 December 1995, G. Monteith, G. Thompson, D. Cook (QM S32915); 1 ♂, Hughes Rd, Topaz, 17°26'S, 145°42'E, pitfall, July–September 1993, G. Monteith, S. Breeden (QM S32911); 1 ♀, Koombooloomba Dam, 4 km S., 17°50'S, 145°36'E, rainforest, pitfall, 9 December 1989–5 January 1990, G. Monteith, G. Thompson, H. Janetzki, (QM S18897); 1 ♀, Maalan SF on H'way, 17°35'S, 145°35'E, rainforest, pitfall, 10 January–31 March 1995, G. Monteith, J. Hasenpusch (QM S32898); 1 ♂, Massey Ck (BS3), 17°37'S, 145°33'E, rainforest, pitfall, 2–30 November




Figure 17 *Amauropelma hasenpuschi* sp. nov.: holotype male (a–e), allotype female (f, g): a, carapace, chelicerae and abdomen, dorsal view; b, leg I, proximal view; c, palpal patella to cymbium, retroventral view showing patella and tibial apophyses; d, e, eyes, dorsal view (d), front view (e); f, g, epigyne, ventral view (f), dorsal view (g). Scale line = 2mm for a, b; 1mm for c–e; 0.3mm for f, g.

1995, L. Umbach (QM S39119); 1 ♂, 1 ♀, Palmerston NP, 17°35'S, 145°41'E, rainforest, pitfall, 25 July–30 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QM S21935, AMNH); 1 ♂, Polly Ck (Hasenpusch property), 17°28'S, 146°00'E, rainforest, intercept flight trap, 7 March–15 May 1995, G. Monteith, J. Hasenpusch (QM S41788); 1 ♂, Stone Ck (Hasenpusch), 17°18'S, 146°00'E, intercept flight trap, 1 October–1 November 1995, J. Hasenpusch (QM S32912); 1 ♂, Tower near The Crater NP, 17°27'S, 145°28'E, rainforest, pitfall, 7 March–15 May 1995, G. Monteith, J. Hasenpusch (QM S41786); 1 ♂, Tower near The Crater NP, 17°27'S, 145°28'E, flight intercept trap, 25 November 1994–10 January 1995, G. Monteith, J. Hasenpusch (QM S31365); 3 ♂, Wongabel SF, 17°19'S, 145°29'E, rainforest, pitfall, 23 July–26 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QM S22616, WAM T42601, AMNH).

Other Material

Australia: Queensland: 1 ♂, Mt Fisher, Kjellberg Rd, 17°31'S, 145°33'E, rainforest, pitfall, 1 December

Table 7 Leg measurements of *Amauropelma hasenpuschi*.

Holotype male (QMS31362).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.61	2.31	2.15	2.85	0.92
Patella	1.31	1.08	0.96	1.15	0.46
Tibia	2.38	1.85	1.54	2.46	0.46
Metatarsus	1.92	1.77	2.00	2.77	
Tarsus	1.15	1.08	1.23	1.77	1.08
Total	9.37	8.09	7.88	11.00	2.92
Allotype female (QMS31363).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.38	1.84	1.92	2.54	1.00
Patella	1.23	1.08	0.92	1.00	0.46
Tibia	1.85	1.38	1.31	2.23	0.69
Metatarsus	1.42	1.38	1.61	2.38	
Tarsus	0.85	0.85	1.08	1.54	0.69
Total	7.73	6.53	6.84	9.69	2.84




Figure 18 *Amauropelma hasenpuschi* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, tibial apophysis, rotated ventral view; c, embolic plate and conductor, ventral view; d, median apophysis, ventral view; e, patellar apophyses, ventral view.

1993–25 February 1994, J. Hasenpusch (QM S32897).

Diagnosis

Differs from most other species in the subquadrate shape of the epigynal plate and from *A. bluewater* in the relatively larger lateral horns and

the rectanguloid median apophysis with very short 'handle'.

Description

Holotype male (QM S31362)

Carapace 3.23 long, 2.46 wide. Abdomen 2.61,

1.62 wide.

Colour. Carapace light brown with dark bands midlaterally and narrow black marginal band. Chelicerae deep reddish brown; legs orange brown with incomplete dark bands: femora, mid- and distal; patellae distal; tibiae mid- and subdistal. Abdomen dorsally pallid with 3 inverted U dark crescents in posterior half and laterally mottled, ventrally pallid with one pair few light shadows medially. Iridescent green sheen on legs. **Eyes.** AME:ALE:PME:PLE, 9:8:7:10. On low tubercles. ALE set along front edge of PME. **Chelicerae.** Promargin with 2 small teeth; retromargin with 4 spaced large teeth and 1 minute basally. **Sternum.** Margins rounded. **Spines.** As for genus except as noted. I: fe pv1p1 d3r1; me p1. II: fe pv1p1 d3r1; pa 0; ti p1; me p1. III: fe pv1p2d3r3; pa p1r1; ti p2d3r2; me p2+2l r2+l2. IV: fe pv1p2d3r2; pa r1; ti p2d2r2; me p2+2l r2+2l v2.2.2.2. **Palp:** fe p1d1.2; pa 0; ti p2d1; cymbium with 2 subapical spines. **Claws.** Tufts small, dense; not as long as in ♀; 3 teeth on I, 1–2 on IV, and basal onychium evident on tarsi. **Legs.** Trochanteral notches deep. **Palp.** Patellar fold normal, broken, small protrusive lip on retrolateral corner. Tibia with retroventral distal widening bearing short, deeply and equally bipartite apophysis; distal tibia with broad triangular extension proventrally and prodorsally (the larger) flanking broad bluntly rounded probasal cymbial process; retrolateral and distal cymbial inner edge a low to high (basally) flange. Subtegulum locking process a blunt cone, tegular interface barely indented. Median apophysis an open gourd-shaped scoop set just inside O-shaped tegulum. Broad, flared conductor near embolic division and begins as circular opening. Embolic plate distally divided into two apices, distal larger, proximal slender. **Spinnerets.** Apices inverted; ALS ca. 1 diameter apart and twice diameter of PLS.

Allotype female (QM S31363)

Carapace 3.23 long, 2.42 wide. Abdomen 4.00, 3.08 wide.

Colour. Carapace and abdomen almost without pattern; legs banded as in ♂. **Eyes.** AME:ALE:PME:PLE, 9:7:8:7. **Chelicerae.** Promargin with 2 teeth; retromargin with 5. **Spines.** As for genus except as noted. I: fe pv1 strong p1d3. II: fe p1d3. III: fe pv1w p2d3r2; pa p1r1; ti p2d3r2; me p2+2l (lower) r2+2l (lower) v1.2.2.2. IV: fe p2d3 r1; pa 0; ti p2d3 r2; me p2+2l r2+2l v2.2.2. **Palp:** fe p1d2; pa 0 ti p1d3; ta p3. **Claws.** Long thin with 3–6 (I, II) or 1 (III, IV) teeth medially. Tufts small distinct on sessile process I, II or raised plate III, IV. All tarsi including palp with basal mound at 1/5. **Spinnerets.** Inverted but ALS separated by 1 diameter, PMS thicker than PLS. **Epigyne.** Externally almost square rounded plate with a rounded vertical horn laterally. Horns presumably receive median apophysis as they are clearly not

ducts but sclerotised guides. Copulatory duct beside horns lead forward to enlarged head and back laterally to enlarged basal receptaculum with long entally directed fertilisation ducts.

Distribution and Habitat

Known only from rainforest at in the central portion of the Wet Tropics Heritage Area, north-eastern Queensland.

Variation

The tibial apophysis of males from different localities shows slight variation in the relative sizes of the apical prongs but otherwise the morphology is acceptably conspecific.

Etymology

For Mr Jack Hasenpusch, the collector of some of the types.

Amauropelma leo Raven and Stumkat, sp. nov.

Figures 4, 19, Table 8

Material Examined

Holotype

♀, Leo's Ck, Coen, Queensland, Australia, 13°32'S, 143°27'E, 25 July 1976, P. Filewood (QM S32903).

Diagnosis

Differs from all other species in the very reduced sizes of the ALE, PME and PLE.

Description

Holotype female (QM S32903)

Carapace 2.32 long, 1.72 wide. Abdomen 2.20 long, 1.40 wide. Total length 5.

Colour. Carapace and legs orange brown, chelicerae darker; abdomen fawn without pattern. **Carapace.** Long fovea extends as far posterior of posterior declivity as anterior. **Eyes.** Small; AME>PME=PLE =ALE. 3 rows, 2.4.2, second row slightly procurved. From above, front of ALE set

Table 8 Leg measurements of *Amauropelma leo*.

Holotype female (QM S32901).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.69	2.56	2.38	3.00	1.31
Patella	1.56	1.44	1.25	1.44	0.50
Tibia	2.38	2.06	1.63	2.50	0.63
Metatarsus	1.88	1.69	1.88	2.75	
Tarsus	0.75	0.81	1.25	1.25	0.88
Total	9.26	8.56	8.14	10.94	3.32




Figure 19 *Amauropelma leo* sp. nov., holotype female. a, carapace, dorsal view; b, eyes, dorsal view; c, epigyne, ventral view. Scale lines = 0.9mm for a, 0.3mm for b, 0.2mm for c.

just anterior to front of PME; ALE clearly closer to PLE than ALE; PLE clearly in separate row to PME. MOQ narrower behind than in front; from front ALE set above AME but part below and part beside PME. Eye group = 0.55 of head/width; PME = 6% of head width. *Chelicerae*. 2 teeth evident on promargin; 5 large and one small basal teeth on retromargin. *Legs*. Tarsal organ in distal third on I, basal 1/6 on III, IV. No scopula. *Spines*. As for genus except as noted. I: fe pv1p1d3. Rest as for *A. mcilwraith* except: II: fe pv1p2d3r1. III: fe p2d3r3. IV: fe p3d3r2. Palp: fe p1d1.2r1; pa p1; ti p2d2; ta p2.1r1.1. *Epigyne*. A broad plate with well rounded wings and soft curving lateral horns.

Male
Unknown.

Distribution and Habitat
Known only from rainforest at Leo Creek, near Coen, Cape York Peninsula.

Etymology
A noun in apposition taken from the type locality.

Amauropelma mcilwraith
Raven and Stumkat, sp. nov.
Figures 3e, 4, 20, 27c, Table 9

Material Examined

Holotype
♂, 11 km W. by N. of Bald Hill, McIlwraith

Range, Queensland, Australia, 13°44'S, 143°20'E, 520m, 27 June–12 July 1989, T.A. Weir (ANIC).

Diagnosis
Differs from all other species in the extensive unsclerotised area surrounding the median apophysis.

Description
Holotype male (ANIC)
Carapace 2.68 long, 2.12 wide. Abdomen 2.20 long, 1.40 wide. Total length 4.8.
Colour. Carapace yellow brown with light marginal and subcentral shadows. Abdomen dorsally pallid with many small dark spots forming no discernible pattern. Legs yellow brown with dark shadows on distal femora, lateral patellae and proximal and distal tibiae and metatarsi, annuli more evident on III, IV, than on I, II. *Eyes*. AME=ALE<PME=PLE. ALE oval; from front, lower

Table 9 Leg measurements of *Amauropelma mcilwraith*.

Holotype male, ANIC.

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.04	1.92	1.76	2.40	1.00
Patella	1.04	0.96	0.80	1.00	0.52
Tibia	1.88	1.40	1.32	2.00	0.52
Metatarsus	1.40	1.32	1.48	2.32	
Tarsus	0.88	0.72	1.20	1.20	1.04
Total	7.24	6.32	6.20	8.92	3.08




Figure 21 *Amauropelma monteithi* sp. nov.: holotype male (a, b, d–f), allotype female (c, h–j): a, carapace, chelicerae and abdomen, dorsal view; b, d, eyes, dorsal view (d), front view (b); c, spinnerets, ventral view; e, f, palpal patella to cymbium, dorsal view (e), retrolateral view showing patella and tibial apophyses (f); g, sternum, maxillae and labium; h, carapace and chelicerae, dorsal view; i, j, epigyne, ventral view (i), dorsal view (j). Scale line = 1mm for a, g, h; 0.5mm for b–f; 0.35mm for i, j.

pallid, 3 irregularly paired pale areas posteriorly. *Carapace*. Pear-shaped; fovea *ca.* as long as mid-carapace width, extends back to slope. Clypeus very narrow < AME. Fine iridescent sheen on carapace and lateral femora, no hairs evident. Anterior constriction very gradual, eyes confined to anterior half. Light cover of fine brown hairs dorsally and on margins; eye region domed downward from about back of anterior constriction. *Eyes*. AME:ALE:PME:PLE, 8:7:6:6. Two very recurved rows. AME on common tubercle and look to front, side and up. ALE look to front and side, PLE to side, PME up and front. Limited dark pigment around all but AME; AME project forward over clypeus. ALE set beside PME. All eyes of similar size and colour. *Chelicerae*. Promargin 3 small basal teeth; retromargin 4 small but larger teeth. Porrect; fang moderately long, boss distinct, triangular. *Labium*. Longer than wide, anterior margin thickened. *Maxillae*. Longer than wide, cylindrical. *Sternum*. Shield-shaped with cuticular extensions at midcoxae. *Legs*. Cuticle translucent. Scopula absent. Pilosity generally

Table 10 Leg measurements of *Amauropelma monteithi*.

Holotype male (QMS31354).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.61	—	2.31	3.23	1.31
Patella	1.31	—	1.00	1.38	0.54
Tibia	2.69	—	1.85	1.85	0.54
Metatarsus	2.23	—	2.00	3.23	
Tarsus	1.31	—	1.08	1.77	1.00
Total	10.15	8.24	11.46	3.39	

Allotype female (QMS24866).

	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.23	1.85	1.69	2.31	1.00
Patella	1.15	0.85	0.77	0.92	0.46
Tibia	1.85	1.46	1.38	2.23	0.54
Metatarsus	1.38	1.31	1.38	2.23	
Tarsus	0.85	0.69	0.92	1.46	0.61
Total	7.46	6.16	6.14	9.15	2.61




Figure 22 *Amauropelma monteithi* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, patellar apophyses, ventral view; c, median apophysis, ventral view; d, tibial apophysis, ventral view.

sparse. Tarsal rod with globular tip at basal one-fifth on all leg tarsi. Trochanters deeply notched, as wide as long. Tarsi enlarge distally. *Spines*. As for genus except as noted. I: fe pv1p1d3r2w; ti p1r1v2.2.2.2; me p1r1. II: fe p1d3 r1; ti p1r1v2.2.2.2; me p1r1. III: fe p3d3r3; pa r1; ti p2d3r2; me p4r4v7. IV: fe p4d3r2; pa r1; ti p2d3 r2v2.2.2; me p4r4v9.

Palp: fe p1d2; pa 0; ti p2r1. *Claws*. Long, slender, gently and uniformly curved; small but distinct claw tufts set well below claws, few hairs around claws. Very prominent anvil-like onychium. Claws reflexed back as in Orsolobidae. 4–6 long close teeth on claws. *Palp*. Patella has ventral flange distally and also distally on retro-edge a small

erect cone directed orthogonal to length. Tibia prolaterally flattened, incrassate on dorsal. Tibial apophysis on subdistal retrodorsal mound, apophysis small apically with shallow division into two cones, one very small basal. Cymbium tear-shaped, without dorsal scopula but with conical hirsute uniformly sclerotised apex and subapical spines; large rounded triangular basal process retroventrally—laterally directed; on proventral basal corner another smaller lobe directly back along length of tibia with spoon-like process distal which presses against dorsal of




Figure 23 *Amauropelma monteithi* sp. nov. Female, scanning electron micrographs: a, distal tarsus, paired claws and claw tufts; b, tarsal rod; c, addressed and erect trichobothria on tarsus.

lateral cymbial lobe. Cymbial groove absent but retroventral lobe arises from flange continuing under bulb and dividing and curving to prolateral basal corner of cymbium ending free. *Spinnerets*. ALS c. equal to PLS in length but ALS thicker, PMS cylinders. All apices domed. ALS c.1 diameter apart noticeably. *Trichobothria*. 2 lines on tibia for length; one line for length of metatarsi; 2 rows, all with trichae of normal length, on tarsi.

Allotype female (QM S24866)

As for male except as follows:

Carapace 2.73 long, 1.88 wide. Abdomen 2.57, 1.73 wide.

Eyes. AME:ALE:PME:PLE, 5:4:3:5. *Spines*. As for genus except as noted. I: fe pv1p1d3. II: fe p1d3. III: fe pv1p1d3r2; pa p1r1; ti p2d3r2; me p4d1r3 v2.2.2. IV: fe p2d3r1; pa r1; ti p2d2r2; me p4r5v2.2.2. *Palp*: fe p1d3; pa 0; ti p2d1; ta p3. *Epigyne*. Externally broad plate or shield with wide wings and distomedial extension lateral of which are two small horns (copulatory channel) with ventral aperture. From copulatory fossae, duct passes forward to small marginal lobe and narrow duct passes back under folded edge of sclerotised shield to distal spherical receptaculum with fertilisation duct off basal inner margin.

Distribution and Habitat

Known only from rainforest at Cape Tribulation, north-eastern Queensland.

Etymology

In honour of Dr Geoff Monteith whose intensive and extensive collections greatly enhance studies on rainforests of north-eastern Australia.

Amauropelma mossman

Raven and Stumkat, sp. nov.

Figures 2, 4, 24–26, Table 11

Material Examined

Holotype

♂, Mossman Bluff track, site 2, Queensland, Australia, 16°25'S, 145°19'E, flight intercept trap, 20 December 1989–15 January 1990, G. Monteith, G. Thompson, ANZSES Expedition (QM S31361).

Paratypes

Australia: Queensland: 1 ♂, Mossman Bluff track, Site 2, 16°25'S, 145°19'E, flight intercept trap, 20 December 1989–15 January 1990, G. Monteith, G. Thompson, ANZSES Expedition (QM S32916); 1 ♂, Mossman Bluff track, Site 2, 16°25'S, 145°19'E, flight intercept trap, 20 December 1989–15 January 1990, G. Monteith, G. Thompson, ANZSES Expedition (QMS31361); 1 ♂, Mossman Bluff Track, 5–10 km




Figure 24 *Amauropelma mossman* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, median apophysis, rotated ventral view; c, patellar apophyses, ventral view; d, tibial apophysis, ventral view.

W. Mossman (Site 2), 16°25'S, 145°19'E, flight intercept trap, 16–30 December 1988, G. Monteith, G. Thompson, ANZSES Expedition (QM S41787); 1 ♂, Clacherty Rd, via Julatten, 16°37'S, 145°19'E, rainforest, sieved litter, 11 October 1980, G. Monteith (QM S32895); 1 ♂, Mossman Bluff track (Site 3), 16°25'S, 145°22'E, pitfall, 16–30 December

1988, G. Monteith, G. Thompson, ANZSES Expedition (QM S32894); 1 ♀, Mossman Bluff track (site 3), 16°25'S, 145°22'E, pitfall, 16–30 December 1988, G. Monteith, G. Thompson, ANZSES Expedition (QM S32893); 2 ♀, Mossman Bluff track (Site 3), 16°25'S, 145°22'E, pitfall, 16–30 December 1988, G. Monteith, G. Thompson, ANZSES

Expedition (QM S32890); 1 ♀, Mossman Gorge NP (NQ 15), 16°28'S, 145°19'E, 1 November 1991–22 July 1992, P. Lawless, R. Raven, M. Shaw (QM S32892); 1 ♀, Mossman Gorge NP (NQ 15), 16°28'S, 145°19'E, 1 November 1991–22 July 1992, P. Lawless, R. Raven, M. Shaw (QM S32891).

Diagnosis

Males differ from those of *A. trueloves* in the size of the retrobasal cymbial process and that the median apophysis is horizontal rather than vertical and from those of *A. hasenpuschi* in the dominant 'handle' on the median apophysis. Females differ from those of *A. wallaman* by the PME clearly being larger than the other eyes.

Description

Holotype male (QM S31361)

Carapace 2.38 long, 1.85 wide. Abdomen 2.00, 1.38 wide.

Colour. Carapace yellowish with wide irregular brown margin and wide brown lateral bands from anterior margins of carapace onto caput and then bracketing fovea; chelicerae yellow brown with dark medial band down each. Abdomen dorsally mottled brown and white with large areas of white mottling anteriorly, ventrally pallid with small dark mottling laterally and posteriorly. Legs yellow brown; femora with brown areas ventrally in undulating waves laterally, two darker bands on tibiae. *Carapace.* Margins hirsute. *Eyes.* AME set close on common tubercle. From above, ALE set

Table 11 Leg measurements of *Amauropelma mossman*.

Holotype male (QM S31361).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.92	1.60	1.52	2.12	0.84
Patella	0.92	0.84	0.72	0.84	0.40
Tibia	1.72	1.32	1.20	1.72	0.28
Metatarsus	1.36	1.24	1.32	2.00	
Tarsus	0.72	0.68	1.20	1.20	0.84
Total	6.64	5.68	5.52	7.88	2.36
Allotype female (QM S32893).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.68	1.44	1.24	1.76	0.60
Patella	0.80	0.80	0.72	0.84	0.40
Tibia	1.36	1.08	1.00	1.64	0.40
Metatarsus	0.96	0.96	1.04	2.00	
Tarsus	0.48	0.40	0.92	0.92	0.56
Total	5.28	4.68	4.72	7.16	1.96

behind anterior edge of PME, tubercle of ALE and PLE touch. AME:ALE:PME:PLE, 7:6:8:6. *Spines.* As for genus except as follows. I: fe pv1p1d3r1. II: fe pv1p1d3r1. III: fe p1p2d3r3. IV: fe pv1p2d3r2. *Palp.* Patella with one distinct cone, ventral corner of fold not enlarged. Tibia with retrolateral apophysis with wide base, apically bifid with short similar apices. Cymbium retrobasally with rounded rectangular lobe, probasal lobe small, rounded; two spines




Figure 25 *Amauropelma mossman* sp. nov., a, b, holotype male: eyes, dorsal view (a), frontal view (b); c, d, allotype female, epigyne, dorsal view (c), ventral view (d). Scale line = 0.3mm.




Figure 26 *Amauropelma mossman* sp. nov. Female, scanning electron micrographs: a, distal tarsus, paired claws and claw tufts; b, tarsal organ; c, adpressed and erect trichobothria.

apico-ventrally on cone. Embolic plate short, beak-like with small foliform conductor; median apophysis set on ventral edge of low C-shaped tegulum, long distinct but tapering handle, lumen subcircular.

Allotype female (QM S32893)

Carapace 2.40 long, 1.76 wide. Abdomen 3.00 long, 2.00 wide. Total length 5.2.

Colour. Carapace yellow with dark narrow marginal and subcentral bands. Abdomen dorsally light green brown with large white mottling. Femora yellow with pale dark finger spots basally, and incomplete band subdistally and distally. **Carapace.** Fovea high, with saddle anteriorly. **Eyes.** 2.4.2. PME=AME>AME=PLE=ALE. ALE-PLE <ALE-AME. From above ALE set behind PME. Back eyes in two well separated rows. ALE look down and to side. From front, lower edge of ALE above upper edge of AME. Eye group = 0.7 of head width; PME = 12% of head width. **Legs.** Tarsi narrow for basal 1/6, widening just distal of tarsal organ at 1/5. Low but large, distinct tarsal organ in basal 1/3 on I; more posteriorly on legs II-IV. All trochanters deeply notched. **Spines.** As for genus except as noted. Tibia I, II with large paired spines overlapping and on raised bases, 4 long predistally and 1 shorter distal; metatarsi with same but only 3 pairs. Pair of strong erect spines proventrally on femur I. I: fe p1pv2d3. II: fe p1d3. III: fe pv1p2d3r1; pa r1; ti p2d2r2v1.2.2; me p3d1r3v2.2.2. IV: fe p2d3r1; pa r1; ti p2d2r2v1.2.2; me p3d1r3v2.2.2. **Palp:** fe d1.2; pa p1; ti p2d2; ta p1.2. **Epigyne.** Plate about 1.5 times wider than long; anterior edge deeply U-shaped; basolateral edge concave with narrow distal edge; large rounded lateral wings; lateral horns fleshy, set anteriorly and wide of wings. **Spinnerets.** ALS distinctly separated.

Distribution and Habitat

Known only from montane rainforest west of Mossman, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

***Amauropelma pineck* Raven and Stumkat, sp. nov.**

Figures 4, 27a, b, d, 28d, Table 12

Material Examined

Holotype

♂, Pine Ck, CSIRO Tower, Queensland, Australia, 17°00'S, 145°50'E, pitfall and intercept traps, 12 September 1991–20 October 1991, G. Monteith, H. Janetzki (QMS 32906).

Paratypes

Australia: Queensland: 4 ♀, same data (QM S39333); 1 ♀, Crystal Cascades, 16°57'S, 145°40'E, pitfall, 27 October 1991–23 July 1992, R. Raven, P. Lawless, M. Shaw (QM S24529); 1 ♀, same data except 23 July 1992–26 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QM S24016); 3 ♀, same data except 26 November 1992–16 April 1993, R., J., and S. Raven, P. and E. Lawless (QM S24882, 24880, WAM T42602). Mission Beach (S1), 17°52'S,

146°03'E, rainforest, pitfall, 29 July–2 September 1996, M. Cermak: 2 ♂ (QM S39103); 1 ♀ (QM S41791); 1 ♀ (QM S41792); 1 ♂ (QM S39102 now WAM T41616); 3 ♂ (QM S41790); 1 ♀, same but 1–19 July 1996 (QM S41789).

Diagnosis

Differs from *A. trueloves* having only four pairs of spines ventrally on tibiae I, II.

Description

Holotype male (QM S32906)

Carapace 2.52 long, 1.92 wide. Abdomen 1.84 long, 1.32 wide. Total length 4.6.

Colour. Carapace yellow brown with narrow dark marginal bands and wide dark bands subcentrally. Abdomen dorsally light greenish brown with large paired four irregular pallid marks. Legs yellow brown with dark shadows medially and distally on femora, laterally on patellae and basally and distally on tibiae, especially on III, IV. *Eyes.* 2.4.2. AME set

Table 12 Leg measurements of *Amauropelma pineck*.

Holotype male (QM S32906).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.80	1.76	1.72	2.24	0.80
Patella	0.92	0.84	0.80	0.96	0.40
Tibia	1.88	1.40	1.16	1.88	0.32
Metatarsus	1.44	1.28	1.36	2.28	
Tarsus	0.72	0.64	1.04	1.04	0.80
Total	6.76	5.92	5.80	8.40	2.32
Allotype female (QM S39333).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.00	1.80	1.68	2.40	0.84
Patella	1.04	0.96	0.80	1.24	0.60
Tibia	1.76	1.40	1.20	1.90	0.44
Metatarsus	1.24	1.20	1.40	2.16	
Tarsus	0.56	0.56	1.16	1.16	0.68
Total	6.60	5.92	5.84	8.86	2.56




Figure 27 *Amauropelma* spp. nov.: scanning electron micrographs. *A. pineck* sp. nov., a, b, d. Male palp, ventral right: a, cymbium and bulb, ventral view; b, distal patella and tibia, retrolateral view; c–g, median apophysis with lumen; c, *A. mcilwraith*; d, *A. pineck*; e, *A. torbjorni*; f, *A. gayundah*, lumen absent; g, *A. rifleck*; f, g, same scale.

forward on common tubercle. MOQ subquadrate. ALE clearly closer to PLE than AME. $PME > PLE = AME > ALE$. From front, lower edge of ALE cut above centres of AME. Back row just forming two rows. *Chelicerae*. 4 teeth on retromargin. *Palp*. Patella without noticeable apophysis. Tibia with retrodorsal conical process with small thorn dorsally at half length. Cymbium ascopulate; prolateral basal lobe squared, retrobasal process rounded, noticeably produced; two spines below cymbial tip. Lumen of subquadrate median apophysis dominant. *Legs*. Tarsal organ at distal third (I), distal 2/5 (II), or basal 1/8 (III, IV). Scopula absent. *Spines*. As for genus except, I: fe pv1d1d3; II, fe pv1p1d2; ti p1. III: fe p2d3r1; pa r1; ti p2d3r2v2.2.2; me p1.1.2r1.1.2.v2.2.2. IV as III but: fe p2d3r2; me p1.1.2r1.2.2.v2.1.1.1.2. Palp: fe d1.1; pa p1; ti p1; cymbium v2.

Allotype female (QM S39333)

Like male except as follows:

Carapace 2.60 long, 2.08 wide. Abdomen 2.60 long, 2.00 wide. Total length 5.9. *Eyes*. Eye group/head width = 0.63; PME = 9% of head width. *Legs*. Patterns darker, distal spines ventrally on tibiae I, II absent. *Spines*. I: fe pv1p1d2; ti v2.2.2.2. II: fe p1d2; ti v2.2.2.2. III: pv1p2d3r1. IV: p1d3r1. Palp: fe d1.2; pa p1; ti p2d1; ta p2.1r.1. *Epigyne*. Two wide lobes or 'wings' with small lateral horns.

Distribution and Habitat

Known only from lowland rainforest near Cairns, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

Amauropelma rifleck Raven and Stumkat, sp. nov.
Figures 3c,d, 4, 27g, 28a–c, 29, Table 13

Material Examined

Holotype

♂, Ellis Beach Rd (NQ 14/2), Queensland, Australia, 16°40'S, 145°34'E, pitfall, 22 July–27 November 1992, R. Raven, P. and E. Lawless, M. Shaw (QMS52203).

Paratypes

Australia: Queensland: 5 ♀, same data (QM S21767, WAM T42603).

Diagnosis

Males differ from those of *A. trueloves* in the large bifid patellar process and median apophysis directed outward, females in the much wider epigynal plates and the ALE are clearly in front of the PME, not beside them as in *A. torbjorni*.

Description

Holotype male (QMS21796)

Carapace 2.68 long, 2.04 wide. Abdomen 1.80 long, 1.32 wide.

Colour. Carapace yellow brown with sheen with darker, wide subcentral band from eyes to posterior slope. Abdomen collapsed away from cuticle. *Carapace*. Very lightly hirsute. *Eyes*. AME on common tubercle. AME:ALE:PME:PLE, 6:5:5:5. From above, ALE set in front of PME; AME on separate mound to PLE. *Legs*. Without tarsal rod. *Spines*. As for genus except as noted. I: fe




Figure 28 *Amauropelma* spp. nov. a–c, *A. rifleck* sp. nov.: holotype male (a), paratype female (QMS 21917 (b, c): a, eyes, dorsal view; b, c, epigyne, ventral view. (b), dorsal view (c); d, *A. pineck* sp. nov., female, epigyne, ventral view; e, *A. wallaman* sp. nov., female, epigyne, ventral view. Scale lines = 0.5mm for a–c, 0.3mm for d, e.




Figure 29 *Amauropelma rifleck* sp. nov.: male palp, ventral right, scanning electron micrographs: a, e, cymbium, bulb and distal tibia, ventral view (a), retrolateral view (e); b, tibia and patella showing apophyses, ventral view; c, patellar apophysis, retroventral view; d, tibial apophysis, retroventral view.

pv1p1d3r1; ti p1r1; me p1r1. II: fe pv1p1d3r1; ti p1r1; me p1r1. III: fe pv1p1d3r2; pa r1; ti p2d3r2v.2.2; me p2r2v.2.2. + distal whorl of 4. IV: fe pv1p2d3r2; pa 0; ti p2d3r2; me p2r2vv.2.2. + distal

whorl of 4. Palp: fe p1d1.2; pa p1r1; cymbium ventroapical 2. *Palp.* Patella with single apically divided retrodistal process apically divided into two cones; tibial apophysis similar but slightly

Table 13 Leg measurements of *Amauropelma rifleck*.

Holotype male.					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.92	1.80	1.72	2.32	1.12
Patella	1.04	0.96	0.84	1.00	0.32
Tibia	1.88	1.52	1.20	2.04	0.40
Metatarsus	1.60	1.40	1.44	1.80	
Tarsus	0.80	0.72	0.80	1.00	0.96
Total	7.24	6.40	6.80	8.16	2.80
Allotype female (QM S21917).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.84	1.72	1.48	2.24	0.88
Patella	1.00	0.92	0.80	0.92	0.44
Tibia	1.60	1.28	1.08	1.88	0.52
Metatarsus	1.20	1.12	1.00	1.92	
Tarsus	0.60	0.56	0.92	0.92	0.72
Total	6.24	5.60	4.96	7.88	2.56

longer. Cymbium without any retrobasal enlargement or lobe. Median apophysis longitudinal, pyriform with lumen not evident in ventral view.

Allotype female (QM S21917)

Carapace 2.44 long, 1.84 wide. Abdomen 2.80 long, 1.48 wide. Total length 5.5.

Eyes. Eye group = 0.66 of head width; PME = 8% of head width. *Spines.* As for male but lacking lateral spines on tibiae and metatarsi I, II *Legs.* Tarsal organ at distal 1/3 on I, distal 3/8 on II, basal on III, IV. *Epigyne.* Plate with wide wings.

Distribution and Habitat

Known only from dense grassy open forest at Ellis Beach, north of Cairns, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

Amauropelma torbjorni Raven and Gray, sp. nov.

Figures 4, 27e, 30, 31, Table 14

Material Examined

Holotype

♂, Keatings Gap, Mt Cook, Queensland, Australia, 15°30'S, 145°13'E, pitfall, 20 July 1992–28 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QMS24364).

Allotype

Australia: Queensland: ♀, Quarantine Rd, Mt

Cook, 15°30'S, 145°13'E, pitfall, 28 November 1992–18 April 1993, R., J. and S. Raven, P. and E. Lawless (QM S24062).

Diagnosis

Males differs from those of *A. trueloves* in larger patellar apophysis; females have a relatively shorter epigyne than those of *A. rifleck* and the ALE are beside PME not in front as in *A. rifleck*.

Description

Holotype male (QM S24364)

Carapace 3.24 long, 2.68 wide.

Colour. Carapace and chelicerae yellow orange with dark lines on interstrial edges. Abdomen collapsed but pallid with dark areas dorsally and mostly pallid ventrally. Legs yellow orange without banding or pattern. *Eyes.* Small; AME set close on common tubercle; from above and in front ALE are beside PME; from in front ALE are transverse and ovoid. AME:ALE:PME:PLE, 6:5:6:4. *Spines.* As for genus except as noted. Very long, strong, on metatarsi especially on III. I: fe p1d2r1; ti p1r1; me p1r1. II: fe p1d4; pa 0; ti p1r1; me p3r2. III: fe pv1p2d4r1; pa r1; ti p2d3r2; me p1.1.2r1.2v2.2.2.2, short pair medially. IV: missing. *Palp.* fe p1d3, pa p1w, ti p2w, cymbium 0 distoventral. *Palp.* Patella with single strong coniform retrodistal process and well separated small ventral prong. Retrolateral tibial apophysis strong, bifurcate with two similar tips; tibia excavate retrolaterally. Strong curved coniform retrobasal process on cymbium and broad rounded probasal process; embolic plate a folded cone with broad translucent conductor, then median apophysis transverse with extensive cavity but overall anvil-shaped with short handle.

Allotype female (QMS 24062)

Like male but:

Spines. As for genus except as noted. I: fe pv1p1d3w. II: fe p1d3w; me v2.2.4. III: fe pv1p2d3r1; pa r1; ti p2d2r2; me p1.1.2d1r1.1.2v2.2.4. IV: fe p1d3r1; pa r1; ti p2d3r2v2.2; me p1.1.2d1r1.1.2v1.1.1.2.2.2. *Palp.* fe p1d3; pa0; ti p2;

Table 14 Leg measurements of *Amauropelma torbjorni*.

Holotype male (QM S24364).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.72	2.44	2.32	—	1.24
Patella	1.40	1.28	1.08	—	0.76
Tibia	2.60	2.00	1.68	—	0.56
Metatarsus	2.04	1.80	1.92	—	
Tarsus	1.00	0.84	0.00	—	1.24
Total	9.76	8.36	7.80	—	3.80




Figure 30 *Amauropelma torbjorni* sp. nov. a, eyes, dorsal view; b, eyes, frontal view; c, d, male palpal patella to cymbium, ventral view (c), retrolateral view (d); e, male carapace, dorsal view; f, epigyne, ventral view. Scales 0.32mm for a, b, 0.25 for rest.




Figure 31 *Amauropelma torbjorni* sp. nov.: male palp, ventral right, scanning electron micrographs: a, cymbium, bulb and distal tibia, ventral view; b, embolic plate, conductor and median apophysis, ventral view; c, palpal tibia and patella showing apophyses, ventral view.

ta v2. *Epigyne*. A broadly rounded plate, wider than long and anteriorly concave, with wide lateral wings and small lateral horns set close to plate.

Distribution and Habitat

Known only from rainforest (type) at Mt Cook, near Cooktown, north-eastern Queensland.

Etymology

Named for Dr Torbjörn Kronestedt, who came to mind while RJR was at the type locality.

Amauropelma undara Raven and Gray, sp. nov.

Figures 4, 32, 33a–c, e–g, Table 15

Material Examined

Holotype

♂, Undara Lava Tubes, Bayliss Cave, Queensland,

Australia, 18°25'S, 144°28'E, 8 February 1996, E. Volschenk, D. Slaney (QM S31389).

Paratypes

Australia: Queensland: same locality as for holotype: 1 ♀, 27 May 1986, F.G. Howarth, D. Irvin (QM S46226); 1 ♂, 1 ♀, 12–13 June 1986, F.G. Howarth, D. Irvin, S. Robson (AM KS50696); 2 ♀, 15 June 1985, F.G. Howarth, D. Irvin, J. Bresnan (AM KS50693, AM KS50694); 1 ♀, 31 May 1986, F.G. Howarth, D. Irvin (AM KS56317); 1 ♂, 14 June 1985, F.G. Howarth, D. Irvin, J. Bresnan (AM KS50692).

Other Material

Australia: Queensland: same locality as for holotype: 1 juvenile, 5 June 1986, F.G. Howarth (QM S46227).




Figure 32 *Amauropelma undara* sp. nov. a, b, male, palpal tibia and tarsus, ventral view (a), with patella (b); c–f, female, c, d, carapace, dorsal view (c), with chelicera, lateral view (d); e, f, epigyne, ventral view (e), dorsal view (f). Scale lines = 0.5mm for a, b, 1mm for c, d, 0.27mm for e, f.




Figure 33 *Amauropelma* spp. *A. undara* sp. nov., a–c, e–g. d, *A. hasenpuschi* sp. nov.: scanning electron micrographs, male. Palp, ventral right: a, cymbium, bulb and distal tibia, ventral view; b, patellar apophysis, rotated retroventral view; c, distal tarsus I, lateral view; d, e, axial view of embolus showing spermatric opening; f, tarsal organ and trichobothrium, dorsal view; g, tarsus showing rugose surface and erect and addressed trichobothria.

Table 15 Leg measurements of *Amauropelma undara*

Holotype male (QMS31389).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	3.38	3.23	2.85	3.85	1.31
Patella	1.31	1.00	1.00	1.00	0.46
Tibia	4.15	3.31	2.85	4.23	0.46
Metatarsus	3.08	2.77	2.61	4.15	
Tarsus	1.61	1.31	1.15	1.77	1.00
Total	13.53	11.62	10.46	15.00	3.23
Allotype female (AMKS50693).					
	Leg 1	Leg 2	Leg 3	Leg 4	
Femur	3.79	3.65	3.42	8.37	
Patella	1.62	1.59	1.40	1.49	
Tibia	3.80	3.55	3.20	4.46	
Metatarsus	2.85	2.85	3.02	4.64	
Tarsus	1.10	1.16	1.22	1.59	
Total	13.16	12.80	12.26	20.55	

Diagnosis

Differs from all other species in the absence of eyes.

Description

Holotype male (QM S31389)

Carapace 2.38 long, 1.85 wide. Abdomen 2.30, 1.58 wide.

Colour. Entirely pallid, only darker cuticle on anterior basal trochanters and palpal bulb. Carapace uniform light cover of moderately long brown bristles. One line of 4–6 bristles arises across on clypeus. Fovea short in a deep groove begins just behind caput ends at edge of back slope. *Eyes.* Entirely absent with low domed mound where AME would be. *Chelicerae.* Promargin with 1 tooth; retromargin with 4 small. Sternum cordate with angles at intercoxae. Maxillae basally with 2 setae flanked by short shallow grooves. *Legs.* Elongate. Iridescent sheen on legs. No scopula, setation sparse. Tarsal organ low, at basal 1/3. Thick spine-

like setae on ventral tarsi. *Spines*. As for genus except as noted. I: fe pv1p1. II: fe p1. III: fe p2d2 r1; pa 0; ti p2d1r2 v0.2.2; me p3r5v2.2.2. IV: fe p1d2 r1; pa 0; ti p2d1r2v2.2.0; me p5r5v2.2.2. *Palp*: fe p1d1.2; pa 0; ti 0. *Claws*. Elongate curved with 3–5 long teeth medially. Tufts small, distinct, no clear division. Trochanteral notches distinct, shallow, 2x wider than deep. *Trichobothria*. One very long basally and medially on metatarsi. *Palp*. Patella with retroventral incision elevated as blunt flap. Tibial apophysis retrodorsal, a narrow process equally divided into 2 flattened triangular points (one broken on ventral right palp in SEM). Cymbium long, with conical apex uniformly setose, prolateral basal corner with blunt darkly sclerotised edge with triangular extension of tibia fitting prodorsal (larger) and proventral (smaller). Embolic plate a short wide grooved process set off tegulum by small translucent hand-shaped conductor. Median apophysis isolated, closed with round aperture. Tegular part of interlocking lobe with distinct point, in front (below) a triangular extension of less sclerotised tegular ring. *Spinnerets*. Short, clearly separated, ALS c.1.5 diameter apart and thicker than PLS but similar length. PMS short cylindrical.

Allotype female (AM KS50693)

Carapace 3.59 long, 2.56 wide. Abdomen 3.12, 2.15 wide.

Colour. Entirely yellowish without darker spots or bands. *Carapace*. Broad, uniformly hirsute; eye region a low mound for about 0.5 of head width; caput long, defined by shallow lateral grooves; fovea short. *Chelicerae*. Large, robust; 4 similar teeth on retromargin, promargin with 3 teeth, middle longest. Dark median zone basally of teeth in large pallid area enclosing basal teeth. *Labium*. Longer than wide, anteriorly indented. *Maxillae*. Longer than wide, about twice length of labium. *Sternum*. Narrowly cordate. *Legs*. Coxae with precoxal sclerites anteriorly and posteriorly. Large pallid retrocoxal hymen on I. All trochanters deeply notched. Preening combs absent. Scopula of long hairs distinct on metatarsi and tarsi I, II; absent on tibiae. Tarsal organ at distal 1/3 on I, II, but at basal 1/4 on III, IV.

Spines. As for genus except as noted. I: fe pv1p2d3r3. II: fe pv1p2d3r3. III: fe pv1p2d3r4; pa p1r1; ti p2d3r2; me p2r2pv1.1rv1.1v2.2.1. IV: fe pv1p2d2r2; pa p1r1; ti p2d3r2; me p3r3v1.1.1.2.2.2. *Palp*: fe fe p1d1.2r1; pa p1r1; ti p2d3r1; ta p2.1.1r3. *Claws*. Long, slender with 3–4 teeth just distal of midpoint. Claw tufts dense, separate, set on (?)erectible bases orthogonal to long tarsal axis. *Spinnerets*. ALS separated; PMS cylinders. *Epigyne*. A broad subquadrate rounded plate with long soft slender lateral

horns; large receptaculum externally evident. An epigynal plug was removed from the anterior ectal corner.

Distribution and Habitat

Known only from caves the Undara lava tunnels via Townsville, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

Amauropelma wallaman
Raven and Stumkat, sp. nov.
Figures 4, 28e, 34, Table 16

Material Examined

Holotype

♂, Wallaman Falls, via Ingham, Queensland, Australia, 18°36'S, 145°47'E, rainforest, sieved litter, 1 October 1980, G. Monteith (QMS 32896).

Allotype

Australia: Queensland: 1 ♀, Cardwell Gap, 18°31'S, 146°11'E, pitfall, 24 September–25 November 1992, R. Raven, P. Lawless, E. Lawless, M. Shaw (QM S24891).

Diagnosis

Males differs from those of *A. gayundah* in that the median apophysis has a dominant lumen evident ventrally. Females differ from those of *A. mossman* in having the PME of similar size to other eyes.

Table 16 Leg measurements of *Amauropelma wallaman*.

Holotype male (QM S32896).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	2.00	1.68	1.84	2.28	1.00
Patella	1.00	1.00	0.84	0.96	0.48
Tibia	1.80	1.28	1.24	1.80	0.48
Metatarsus	1.56	1.40	1.48	2.08	
Tarsus	0.84	0.80	1.28	1.28	1.04
Total	7.20	6.16	6.24	8.40	3.00
Allotype female (QM S24841).					
	Leg 1	Leg 2	Leg 3	Leg 4	Palp
Femur	1.92	1.68	1.60	2.04	0.88
Patella	0.92	0.92	0.80	0.92	0.52
Tibia	1.44	1.32	1.12	1.80	0.52
Metatarsus	1.16	1.08	1.20	1.92	
Tarsus	0.60	0.52	1.00	1.00	0.60
Total	6.04	5.52	5.40	7.68	2.52




Figure 34 *Amauropelma wallaman* sp. nov.: scanning electron micrographs, male. Palp, ventral right: a, b, cymbium, bulb and distal tibia, ventral view (a), retrolateral view (b); c, patellar apophysis, retroventral view; d, tibial apophysis, retrolateral view.

Description

Holotype male (QM S32896)

Carapace 2.44 long, 2.00 wide. Abdomen 2.00, 1.20 long.

Colour. Carapace light orange brown with narrow dark edge, dark lines along striae edges and on caput. Legs collapsed but yellow brown with slightly darker bands distally on femora and basally on tibiae. Abdomen pallid with dark areas laterally and distally forming paired long ovoid zones anteriorly and 3 transverse bars posteriorly. **Carapace.** Glabrous except for fine hairs along margins. **Eyes.** AME on common tubercle. ALE and PLE with bases of tubercle touching. Sinuous, loosely grate-shaped tapetum just evident. All eyes about equal in size. **Spines.** As for *A. rifleck*. **Palp.** Patella with two dissimilar small cones, ca. 1 diameter apart. Tibia with low retrodorsal mound with deeply divided tip; distal tip about 2–3 diameters of basal tip. Cymbium with short but distinct retrobasal process, and large probasal heel; no ventral spines on apical cone. Tegulum C-shaped but basally extends out in broadly rounded 'chin', median apophysis subovate with large extensive lumen and set basally on tegulum, longitudinal and set precariously on edge; conductor broadly flared, embolic plate with folded groove.

Allotype female (QM S24891)

Carapace 2.56 long, 1.84 wide. Abdomen 2.80 long, 1.92 wide. Total length 5.7.

Colour. Carapace yellow brown with dark sashes laterally and dark margins. Abdomen colour lost. Legs yellow brown with mid- and distal bands on femora and tibiae, central bands on patellae and weakly on metatarsi. **Eyes.** Group width/head width = 0.64; PME = 10% of head width. **Spines.** As for genus except as noted. On tibia I, (ventral) spines are distally higher on lateral face than basally. I, II: fe p1d3. III: fe pv1p2d3r1; pa p1r1; ti p2d3r2; me p2d2r2v2.2.2. IV: fe p2d3r1; pa r1; ti p2d2r2; me p2d2r2v2.2.2. **Palp:** fe p1d1.2; pa p1; ti p1d1; ta p3. **Spinnerets.** ALS with slightly enlarged spigot medially on tip. PMS with 2 enlarged spigots. **Epigyne.** A plate about 1.5 times wider than long with short soft lateral horns.

Distribution and Habitat

Known only from rainforest at and near Wallamans Falls, near Ingham, north-eastern Queensland.

Etymology

A noun in apposition taken from the type locality.

ACKNOWLEDGEMENTS

This paper is dedicated to Dr Glenn Hunt, 1944–

1999, colleague, friend and highly esteemed scientist who inspired not only by his professional but also by his personal example of courage and dedication. We are very grateful to Dr Diana Silva, American Museum of Natural History, New York, for provocative discussions, comments on the manuscript and especially the suggestion to compare *Amauropelma* to *Thoriossa* Simon. Drs Barbara Baehr and V. Davies, Queensland Museum, provided support and counterpoint through the final phases. Mrs Helen Stark skillfully illustrated *Amauropelma mossman* (Figure 1) and Dr Barbara Baehr very kindly inked figures of epigynes and commented on the manuscript. Ms Wendy Rawlinson, Queensland Museum, very ably helped with registration and sorting of material. Dr Rudy Jocqué, Musée Royal de l'Afrique Centrale, Tervuren, Belgium, kindly loaned the material of *Thoriossa spinivulva* Simon (RMAC 131.124) and *Thoriossa taurinain* Simon (RMAC 131.101). For all of that, we are very grateful.

REFERENCES

- Benoit, P.L.G. (1976). Etudes sur les Ctenidae africains (Araneae). II. Les genres *Thoriossa* Simon et *Trogloctenus* Lessert. *Revue Zoologiques Africains* 90: 221–227.
- Davies, V. (1976). Spiders. In: *Fauna of Eastern Australian Rainforests: preliminary report on sites surveyed by the Queensland Museum in mid-eastern and north-eastern Queensland*. Queensland Museum, Brisbane.
- Davies, V. (1977). Spiders. In: *Fauna of Eastern Australian Rainforests II*. Queensland Museum, Brisbane.
- Davies, V. (1986). New Australian species of *Otira* Forster and Wilton, 1973 and *Strenosoma* Hogg, 1900 (Araneae, Amaurobiidae). *Memoirs of the Queensland Museum* 22: 237–251.
- Dippenaar-Schoeman, A.S. and R. Jocqué (1997). *African Spiders. An identification manual*. Plant Protection Research Institute Handbook No. 9. Biosystematics Division, ARC-Plant Protection Research Institute, Ultra Litho, Johannesburg.
- Forster, R.R., Wilton, C.L., 1973. The spiders of New Zealand. Part IV. Agelenidae, Stiphiidae, Amphinectidae, Neolanidae, Ctenidae, Psechridae. *Otago Museum Bulletin* 4: 1–309.
- Gray, M.R. (1973). Cavernicolous spiders from the Nullarbor Plain and south-west Australia. *Journal of the Australian Entomological Society* 12: 207–221.
- Griswold, C.E. (1991). A revision and phylogenetic analysis of the spider genus *Machadonia* (Araneae, Lycosoidea). *Entomologia Scandinavica* 22: 305–351.
- Griswold, C.E. (1993). Investigations into the phylogeny of the lycosoid spiders and their kin (Arachnida, Araneae, Lycosoidea). *Smithsonian Contributions to Zoology* 539: 1–39.
- Griswold, C.E. (1994). A revision and phylogenetic analysis of the spider genus *Phanotea* Simon (Araneae, Lycosoidea). *Annales Sciences Zoologique* 273: 3–63.

- Koch, L. (1878). *Die Arachniden Australiens, nach der natur beschrieben und abgebildet*. Vol. 1: 969–1064. Bauer and Raspe, Nürnberg.
- Lehtinen, P.T. (1967). Classification of the cribellate spiders and some allied families, with notes on the evolution of the suborder Araneomorpha. *Annales Zoologici Fennici* 4: 199–467.
- Platnick, N.I. 2000. A relimitation and revision of the Australasian ground spider family Lamponidae (Araneae: Gnaphosoidea). *Bulletin of the American Museum of Natural History* 245: 1–330.
- Rainbow, W.J. (1911). A census of Australian Araneidae. *Records of the Australian Museum* 9: 107–319.
- Raven, R.J. (1981). A review of the Australian genera of the mygalomorph subfamily Diplurinae (Dipluridae: Chelicerata). *Australian Journal of Zoology* 29: 321–363.
- Raven, R.J. (1994). Mygalomorph spiders of the Barychelidae in Australia and the western Pacific. *Memoirs of the Queensland Museum* 35: 291–706.
- Raven, R.J. (1998). Revision of the Australian genera of the Miturgidae with a preview of their relationships. *XIVth International Congress of Arachnology, Chicago, 1998. Abstracts*: 31.
- Thorell, T. (1881). Studi sui Ragni Malesi e Papuani. III. Ragni dell'Austro-Malesia o del Capo York, conservati nel Museo Civico di Storia Naturale di Genova. *Annali del Museo Civico di Storia Naturale Giacomo Doria, Genova* 17: 1–720.

Manuscript received 27 April 2000; accepted 21 December 2000.